



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

Propolis: A review of its biological actions and health benefits.

Propolis, a review of its biological actions and health benefits

Propóleo, a review of its biological actions and benefits for health

Edson Geronimo, Pharmacy student at the University of Paraná (UNIPAR), PhD professor at UNIPAR in health-related courses, Umuarama/PR

João Vitor Barbosa do Carmo, Pharmacy student at the University of Paraná (UNIPAR), PhD professor at UNIPAR in health sciences courses, Umuarama/PR

Juliana Cogo, PhD Lecturer in health science courses at the Cesumar University Center (UNICESUMAR), Umuarama/PR.

Eliane Mendes Franco, students of the Professional Master's Program in Medicinal Plants and Phytotherapy in Primary Care at the Paranaense University (UNIPAR), Umuarama/PR.

Simony Rodrigues Bernardelli Rosa, students of the Professional Master's Program in Medicinal Plants and Phytotherapy in Primary Care at the Paranaense University (UNIPAR), Umuarama/PR.

Rosinéia Aparecida Vilela Cebrian, students of the Professional Master's Program in Medicinal Plants and Phytotherapy in Primary Care at the Paranaense University (UNIPAR), Umuarama/PR.

Ezilda Jacomassi, PhD Professor in the Professional Master's Program in Medicinal Plants and Phytotherapy in Primary Care at the Paranaense University (UNIPAR), Umuarama/PR.

Emerson Luiz Botelho Lourenço, PhD Professor in the Professional Master's Program in Medicinal Plants and Phytotherapy in Primary Care at the Paranaense University (UNIPAR), Umuarama/PR.

Daniela de Cassia Faglioni Boleta-Ceranto, PhD Professor in the Professional Master's Program in Medicinal Plants and Phytotherapy in Primary Care at the Paranaense University (UNIPAR), Umuarama/PR.

Giuliana Zardeto, PhD, Professor at the University of Paraná (UNIPAR) in health-related courses, supervisor of final course projects, Umuarama/PR.

SUMMARY

Propolis is a mixture of varying amounts of beeswax and resins collected by bees from plants, particularly buds and resinous exudates. The composition of propolis varies according to the type of bee, geographic origin, and plant source of the samples. Propolis is important for the defense of the hive and has been used for its medicinal properties since antiquity. The biological characteristics of propolis depend on its chemical composition, plant sources, geographic zone, and seasons. More than 300 compounds have been identified in propolis, including phenolic compounds, aromatic acids, essential oils, waxes, and amino acids. Many scientific articles are published every year in different international journals, and several research groups have focused their attention on the chemical compounds and biological activity of propolis.

Keywords: Propolis, Pharmacological activities, Research.

ABSTRACT

Propolis is a mixture of varying amounts of beeswax and resins collected by bees from plants, particularly buds and resinous exudates. The composition of propolis varies according to the type of bee, geographical origin and plant origin of the samples. Propolis is important for the defense of the hive and has been used for its medicinal properties since ancient times. The biological characteristics of propolis depend on its chemical composition, plant sources, geographical area and seasons. More than 300 compounds have been identified in propolis, such as phenolic compounds, aromatic acids, essential oils, waxes and amino acids. Many scientific articles are published every year in different international journals, and several groups of researchers have focused their attention on the chemical compounds and biological activity of propolis.

Keywords: Propolis, Pharmacological activities, Research.

ABSTRACT

The propoleum is a mixture of variable amounts of beeswax and resins collected by bees from plants, particularly from buds and resinous exudates. The composition of the oil varies depending on the type of bee, and the geographic and vegetal origin of the samples. The propellant is



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

It is important for the defense of the hive and has been used for its medicinal properties since ancient times. The biological characteristics of the oil depend on its chemical composition, plant sources, geographic zone and seasons. More than 300 compounds have been identified in the oil, such as phenolic compounds, aromatic acids, essential oils, waxes and amino acids. Each year numerous scientific articles are published in different international journals and several groups of researchers focus their attention on chemical compounds and the biological activity of the oil.

Keywords: Propolis, Pharmacological activities, Research.

INTRODUCTION

The term propolis derives from the Greek pro (para, 'in front of', 'at the entrance of') and polis. ('community' or 'city') and means a substance in defense of the hive (CASTALDO; CAPASSO, 2002). Propolis is a resinous substance collected by *Apis mellifera* from various buds of trees that they use to line parts of the hive and seal cracks and crevices in the hive (TORETI et al., 2013). Propolis has been used as a folk medicine since 300 BC (SUNG et al., 2017).

Propolis was used as an antiseptic and healing agent in the treatment of wounds and as mouth disinfectant, with these uses being perpetuated in the Middle Ages and among Arab physicians. Propolis was also recognized by other peoples unrelated to the civilizations of the Old World: The Incas used propolis as an antipyretic agent, and the London pharmacopoeias of the 17th century... They listed propolis as an official drug. Between the 17th and 20th centuries, propolis became very... popular in Europe due to its antibacterial activity (CASTALDO; CAPASSO, 2002).

The medical application of propolis preparations has led to an increased interest in their use. chemical compositions and their botanical origins, because until now mainly compounds Polyphenolics were identified in propolis collected by *Apis mellifera*. Flavonoids, the The main polyphenols in propolis vary quantitatively or qualitatively, depending on the environmental ecology of the plant (TORETI et al., 2013).

Recently, numerous biological properties of propolis have been reported, including cytotoxic, anti-herpes, free radical scavenging, antimicrobial and anti-HIV. Due to its wide range of biological activities, propolis has been widely used in Foods and beverages to improve health and prevent disease (RAO et al., 2016). However, the The chemical composition of propolis is susceptible to geographic location, origin, and bee species. (SALATINO et al., 2011; TORETI et al., 2013). The objective of this literature review was to report the Biological and functional properties of propolis produced by bees.

2. MATERIALS AND METHODS

This study aimed to conduct a literature review using the Google Scholar® database for the period 1967 to 2025. The keywords used were: extract



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

The role of propolis in boosting immunity and overall health: understanding its main effective active ingredients. against microorganisms, acting as a natural immunizing agent due to its active ingredients, being recognized as a bioactive supplement.

3. LITERATURE REVIEW

3.1 Propolis

Propolis is a natural resinous mixture produced by bees from substances collected from plant parts, buds, and exudates. The word propolis is derived from Greek, where "Polis" means "at the entrance of" and "polis" means "community" or "city," which means that this product Natural is used in the defense of the beehive (WHAG, 2013).

As defined by Lewis and Short, propolis is considered the third product in manufacturing of honey, described as a gummy substance with which bees seal the cracks in their sacs. Beehives, also called "bee glue." It is generally believed that bees produce Propolis helps protect the beehive. In addition to its role in sealing holes and blocking cracks... and soften the inner walls, bee glue appears to act as an antiseptic to prevent... Microbial infection of larvae, honey deposits, and combs (WAGH, 2013). As Breyer et al. (2016) They observed that bees apply propolis to the areas where the honeycombs are to be attached, creating surfaces. smooth and germ-free.

Because bee populations are so confined and live in close contact, the disease... In one bee, the infection can spread rapidly throughout the entire hive. However, the hives They remain healthy because bees manufacture their own antibiotic (DIDARAS et al., 2020). In other words, propolis, which reduces microbial growth on the walls of beehives. Furthermore, the Propolis protects the beehive against uncontrolled airflow and external moisture. The thin layer of Propolis provides a waterproof lining that limits water leakage and maintains constant moisture in the interior of the beehive (SIHERI et al., 2017).

3.2 Chemical composition of propolis

Propolis is the third most important component of bee products. It is composed of... mainly composed of resin (50%), wax (30%), essential oils (10%), pollen (5%) and other organic compounds (5%) (GÓMEZ-CARAVACA, 2006). Phenolic compounds, esters, flavonoids, terpenes In oxidative medicine and cellular longevity, beta-steroids, aromatic aldehydes, and alcohols are the important organic compounds present in propolis (HUANG et al., 2014).

Twelve different flavonoids, namely pinocembrin, acacetin, chrysin, rutin, luteolin, kaempferol, apigenin, myricetin, catechin, naringenin, galangin and quercetin; two phenolic acids, caffeic acid and cinnamic acid; and a stilbene derivative called resveratrol. They were detected in propolis extracts by capillary zone electrophoresis (VOLPI, 2004).



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

Propolis also contains important vitamins, such as vitamins B1, B2, B6, C and E.

Useful minerals such as magnesium (Mg), calcium (Ca), potassium (K), sodium (Na), copper (Cu), zinc (Zn), manganese (Mn) and iron (Fe). Some enzymes, such as succinate dehydrogenase, glucose-6-phosphatase, Adenosine triphosphatase and acid phosphatase are also present in propolis (LOTIFY, 2006).

3.2.1 Variations in the chemical composition of propolis

Propolis is a resinous mixture made by bees from substances collected from tree buds or other plants, plant exudates or resins found on the stem, branches or leaves of different plants. The geographical origin of propolis is given by the plant sources of respective areas. Different studies have classified this bee product according to the material. Plants from the same areas. The physical appearance of propolis varies greatly depending on its origin. The botanical and geographical characteristics, as well as the type of bee involved in the collection process, influence the color of propolis. cream, yellow, green, light brown, dark brown or red (DELAPLANE, 2010).

The chemical composition of propolis varies greatly, depending mainly on seasonality. locality and vegetation (VALENCIA et al. 2012). The composition of propolis varies not only from one Differences can occur between different types, but even between samples of the same type. For example, differences have been observed when comparing the composition of red propolis from Cuba and Brazil (GIMÉNEZ-CASSINA et al.). 2014).

Dalbergia ecastaphyllum is recognized as the plant source of resin for propolis. Brazilian and Cuban red propolis. However, only Brazilian red propolis has been reported as possessing a second plant source, probably a species of *Clusia*, from which they derive. prenylated benzophenones, namely gutiferone E, xanthokimol and oblongifolin A (PICCINELLI et al. (2011). The effect of seasonality on the composition of Brazilian propolis has been extensively studied. studied (SOUZA et al. 2016; BUENO-SILVA et al. 2017; REGUEIRA NETO et al. 2017). Seasonal quantitative chemical differences have also been reported for red propolis. Brazilian (BUENO-SILVA et al. 2017).

Several factors can explain the seasonal variations in propolis. The composition of the oils. The availability of essential plants is susceptible to seasonal factors (HUSSAIN et al. 2010). The plant parts that provide resin vary according to the season, especially in certain species. Annual or biennial. Bees depend on the availability of young tissues of *B. dracunculifolia*. for the production of propolis (TEIXEIRA et al. 2005). During the plant's reproductive period. (December-May), little or no young vegetative tissue is available (BASTOS et al. 2011). With regard to *Clusia fluminensis* species, the opposite occurs, as the resin they provide and The substances used by bees for the production of propolis are exudates from female flowers (CUESTA-RUBIO et al. 2002).

The main source of Brazilian propolis is the leaf resin of *Baccharis dracunculifolia*.



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025
including constituents such as diterpenes, lignans, prenylated p-coumaric acid derivatives such as
as well as acetophenone and flavonoids. Brazilian propolis contains a large amount of artepillin C.
quantity when compared with caffeic acid phenethyl ester (CAPE) (CHAN et al., 2013;
FERREIRA et al., 2017).

3.3 Bioactive compounds of propolis

Propolis is highly rich in bioactive compounds. Essential and non-essential compounds,
Polyphenols and vitamins, which occur naturally as part of food chains, are
considered bioactive. These compounds are naturally present in foods and confer
Phenolic compounds are bioactive compounds with beneficial health properties. They are defined as:
as organic compounds with an aromatic ring that is chemically bonded to one or more
hydrogenated substituents in the presence of corresponding functional derivatives (MARIN et al.,
2002).

In propolis, phenolic compounds are commonly present as flavonoids (KÜÇÜK
et al., 2007). Several phenolic compounds contribute to the functional properties of the products.
apicultural, including their antioxidant, antimicrobial, antiviral, and anti-inflammatory activities,
antifungal, wound-healing and cardioprotective (BIESALSKI et al., 2009).

3.4 Functional activities of propolis

Depending on its chemical composition, propolis has antimicrobial properties.
(antibacterial, antifungal, antiviral), antioxidant, anti-inflammatory, immunomodulatory,
Antitumor, antiulcer, hepatoprotective, and cardioprotective properties. Due to its composition and
Due to its multifactorial activities, propolis has been suggested to show potential benefits for human health;
the biological activities of propolis depend on its constituents (Küçük et al.,
2007; BIESALSKI, 2009; CHAN et al., 2013).

3.4.1 Antimicrobial activity

Propolis exerts antimicrobial activity against a wide range of microorganisms.
such as bacteria, fungi and viruses (HEGAZI, 2002; ORSI et al., 2005; VECCHI; DRAGO, 2007;
NOLKEMPER et al., 2009; POPOVA et al., 2009; SHNILTZLER et al., 2009; SHIMIZU et al.,
(2009). The fractions of Brazilian propolis have an efficient antibacterial action mainly
against Gram-positive bacteria (POBIEGA ET AL., 2019; BARREIRAS et al., 2020).

The formation of tooth decay is caused by the colonization and accumulation of oral microorganisms.
and extracellular polysaccharides that are synthesized from sucrose by glycosyltransferase of
Streptococcus mutans (AKCA et al., 2016). The ethanolic extract of propolis, containing the largest
Concentrations of pinocembrin and galangin have been shown to cause marked inhibition of the activity of
glycosyltransferase and reduced growth of S. mutans, implying its potential use in prevention.



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025
of dental caries and oral diseases (DE LUCA et al., 2017).

Furthermore, Salmonella spp. infection is a major health problem worldwide. worldwide, causing diarrhea, fever, and abdominal cramps. High concentration of ethanolic extract of Propolis significantly inhibits the growth of Salmonella, suggesting the possible use of propolis as an alternative for controlling Salmonella infection. Its stronger antibacterial activity. The limited effectiveness against Gram-positive bacteria and its limited action against Gram-negative bacteria may be due to... richness in flavonoids (SOUZA et al., 2013; AL-ANI et al., 2018).

3.4.2 Antiviral activity

Both aqueous and ethanolic extracts of propolis exhibit high antiviral activity. Although aqueous and ethanolic extracts of propolis contain different polyphenols, flavonoids, and acids. Phenylcarboxylic acids, aqueous propolis. The extract contains a relatively high amount of acids. Phenylcarboxyls and a low concentration of flavonoids compared to the ethanolic extract. Both propolis extracts exhibit high levels of antiviral activity against HSV-1 and HSV-2. Different components of propolis in its extract exhibit a greater anti-herpetic effect and greater selectivity than the single constituent, therefore topical application of propolis may be suitable against herpes infection (SCHNITZLER et al., 2010; LABSKÁ et al., 2016).

3.4.3 Antioxidant activity

Propolis has been observed to possess antioxidant properties due to its components. galangin and pinocembrin (EL-GUENDOUC ET AL., 2017; MACHADO et al., 2017). Due to Due to its higher polyphenol content, the aqueous extract of propolis was more effective than the ethanolic extracts. Galangin showed greater activity in both extracts compared to pinocembrin. due to the structural difference in both (YINKANG et al., 2018; NICHITOI, 2021).

Antioxidants have the ability to deactivate free radicals and also prevent the... Vitamin C, lipids, and other compounds were destroyed or oxidized. Because free radicals and other factors are the main cause of cell aging and deterioration in states such as Parkinson's disease, Alzheimer's disease, arthritis, cancer, diabetes, cardiovascular diseases (SHARIFI-RAD et al., 2020) and impaired liver function. Components of propolis such as vanillin Phenolic acids have the ability to penetrate the epidermis and dermis and protect them from free radicals. free cells, produced due to radiation or before the maturation of dermal cells due to aging. (ANJUM et al., 2019).

The antioxidant properties of propolis are due to the phenolic compounds that... They donate hydrogen ions to free radicals to protect cells from oxidation reactions and also Propolis has the ability to protect food from oxidation and poisoning. To remove free radicals, which are the main cause of lipid, acid, and protein oxidation.



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025 (KOKOT et al., 2018). The extract of Portuguese propolis demonstrated an antioxidant property and to prevent lipid peroxidation in human red blood cells (ZABAIUO et al., 2017).

3.4.4 Anti-inflammatory activity

The anti-inflammatory activity of propolis not only suppresses the generation of prostaglandins and leukotrienes inhibiting the expression and activities of cyclooxygenases and lipoxygenases, but also It delays the gene expression of nitric oxide synthesis, and the activation of NF- κ B mediated by TNF- γ and reducing the immune response in T cells (FRANCHIN et al., 2016; EL-GUENDOZ et al. 2017). THE Inhibition of NF- κ B activation may be the molecular basis for its anti-inflammatory properties. of propolis (XUAN et al., 2019).

Woo et al. (2005) investigated the effect of chrysin on the expression of cyclooxygenase-2 (COX-2) in lipopolysaccharide (LPS)-activated Raw 264.7 cells. They reported that chrysin suppresses significantly the LPS-induced COX-2 protein and mRNA expression in a way Dose-dependent. Mutation analysis and electrophoretic mobility change assay. They found that the nuclear factor for IL-6 is responsible for the COX-2-mediated downregulation. by chrysin, suggesting its anti-inflammatory and anticancer properties (WOO et al., 2005; YOSEFI et al., 2022).

3.4.5. Immunomodulatory activity

The immunomodulatory action of propolis is well studied, but the exact mechanisms of The effects of propolis on the immune system are still unclear. In vitro and in vivo assays show that the action The immunomodulatory effect of propolis on murine peritoneal macrophages is not caused solely by increased microbicidal activity, but also due to the increased stimulating action on lytic activity. of natural killer cells against tumor cells (SFORCIN, 2007; FISCHER et al., 2008). You Ethanol extracts of propolis stimulate antibody production in immunized mice. with sheep red blood cells (AL-HARIRI, 2019). The enzyme, indoleamine 2,3-dioxygenase, regulates the immune responses through the ability to degrade tryptophan into its metabolites, which are responsible for suppressing the function of effector T cells and promoting the differentiation of T cells Regulators. Increasing the dose of ethanol extract of propolis or increasing the number of its Administration of splenic cells from immunized mice produces an inhibitory effect on formation of arterial plaques (ALMEIDA; MENEZES, 2002; PARK et al., 2008).

3.4.6. Hepatoprotective activity

Several studies have reported the hepatoprotective effect of propolis in liver injuries. Animal models of hepatotoxicity can be created by inducing liver damage with administration of Paracetamol (WILLIAMS et al., 2011), or administering Concanavalin A (WANG et al., 2012). Ye et al. (2019) induced chronic alcohol-induced liver damage in rats.



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

adding alcohol to their daily diet for seven weeks. This group administered different doses of Propolis (ethanolic extract, three times a day for three days, orally) after seven weeks of ingestion of alcohol. The ethanolic extract of propolis (EEP), the therapeutic effects of EEP may be mediated by modulation of metabolic processes such as lipid biosynthesis and fatty acid oxidation. These effects associated with decreased plasma levels of ALT, ALP, and TC contributed to to improve alcohol-induced fatty liver (YE et al., 2019). As mentioned above, the The chemical composition of propolis varies considerably depending on its geographic origins and sources. vegetables.

Banskota et al. (2000) investigated three different biological activities (activity of elimination of DPPH free radicals, cytotoxicity and hepatoprotective activity) in methanol, as well as in aqueous extracts of propolis from Brazil, Peru, the Netherlands, and China. The elimination activity Stronger DPPH free radical scavengers are found in aqueous extracts of six Brazilian propolis varieties and Chinese extracts are more potent than their corresponding methanolic extracts, while the methanolic extracts of Dutch and Peruvian propolis have stronger DPPH free radical scavenging activity. than their corresponding aqueous extracts.

However, the methanolic extracts of all propolis samples in this study exhibit stronger cytotoxicity than the corresponding aqueous extracts; while almost all of the The samples have significant hepatoprotective activity (BANSKOTA et al., 2000). The activity The hepatoprotective effect of the alcoholic extract of Brazilian tropical propolis is mainly due to Phenolic compounds, including flavonoids. In the same experimental model, diterpenes of the type Labdane (isolated from the methanolic extract) also possesses significant hepatoprotective activity. (BANSKOTA, et al., 2001).

Collective evidence suggests that propolis has a marked hepatoprotective potential. due to its composition of flavonoids, phenolic compounds, and diterpenes.

3.4.7 Cardioprotective activity

Flavonoids act as natural antioxidants and iron chelators, therefore they have been... used as cardioprotective agents in doxorubicin-induced cardiotoxicity caused by the formation of oxygen free radicals (KAISEROVÁ et al., 2007). However, the mechanism of action The underlying cause of all the flavonoids present in propolis extract remains elusive.

Pre-treatment of these rats with propolis extract administered 4 days prior to Doxorubicin and/or vinblastine not only substantially reduced peroxidative damage in tissues. myocardial, but also restored catalase and SOD activities, supporting the view that the Polyphenols in propolis protect heart tissue by blocking oxidative stress and restoring mitochondrial dysfunction (ALYANE et al., 2006). ROS generation favors the release



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

of cytochrome c from mitochondria and consequent formation of the apoptosome complex through Positive regulation of Bax or negative regulation of Bcl-XL inducing apoptosis. CAPE exerts an effect. antiarrhythmic (reduces the incidence and duration of ventricular tachycardia and ventricular fibrillation, as well as such as decreased mortality) in rats subjected to myocardial ischemia and ischemic injury- Reperfusion. The ethanolic extract of propolis at a dilution of 10⁻⁵ exhibits the highest elimination activity. of free radicals when compared with vitamin E and quercetin (HUANG et al., 2005).

These observations support the view that some of the protective effects of propolis may to be attributed to the direct elimination of CAPE properties (HUANG et al., 2014). In addition, the Administration of propolis in diabetic rats decreases blood glucose levels, malondialdehyde, NO, NOS, total cholesterol, triglycerides, low-density lipoprotein cholesterol, cholesterol of very low-density lipoprotein in the serum of fasting rats and increases serum levels of high lipoprotein density, cholesterol, and superoxide dismutase, suggesting that propolis regulates the The metabolism of glucose and lipids in the blood leads to a decrease in the production of Lipid peroxidation and free radical scavenging in rats with diabetes mellitus (KHALIL, 2006).

3.4.8 Anticancer activity

Due to its antioxidant activity, the ethanolic extract of propolis (EEP) taken from Indian stingless bee colonies showed anticancer properties against four cancer cell lines at various concentrations, which resulted in apoptosis and cytotoxicity of these cancer cells (KRÓL et al., 2013).

The flavonoids in propolis have been shown to halt breast cancer, lung cancer, and oral cancer. as well as esophageal, stomach, colorectal, prostate and skin cancer (MARTINOTTI; RANZATO, (2015). Brazilian propolis has angiogenic properties, in addition to preventing an increase in the number of human umbilical vein endothelial cells (ZABAIU et al., 2017). The ethanolic extract of Brazilian propolis showed anticancer properties when tested on 1,2-Dimethylhydrazine. which causes colon carcinogenesis in mice (WATANABE et al., 2011). The aqueous extract Propolis from Thailand, examined by researchers, showed greater anticancer activity. against the SW620 colon carcinoma cell line compared to the methanolic extract of propolis (WATANABE et al., 2011).

The ethanolic extract of propolis showed cytotoxicity against adenocarcinoma cells. HT-29 colonoscopy, as well as HT-1080 human fibrosarcoma, but did not demonstrate cytotoxicity to typical human skin fibroblasts (WATANABE et al., 2011).

3.4.9 Antitumor activity

The components of propolis have antitumor properties (KOMERICKI; KRÄNKE, 2009; VEIGA et al., 2017) and components such as caffeic acid phenethyl ester (CASTALDO;



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

CAPASSO (2002) and artepillin C were investigated and shown to possess antitumor effects.

(CHAN et al., 2013). These propolis compounds are involved in cell cycle arrest,

inhibition of matrix metalloproteinases, anti-angiogenic effect, and also inhibit the transfer of

Diseases spread from one part of the body to another (SFORCIN, 2016). Propolis has the ability to...

Interrupting DNA synthesis in tumor cells has the property of causing aging.

of tumor cells (apoptosis) and has the ability to trigger white blood cells to generate

those agents capable of regulating the function of B, T, and natural killer cells, respectively (SALOMÃO et al.,

2011; WAGH, 2013). Other compounds such as galangin, cardanol, nemorosone

Chrysin and other enzymes are involved in preventing the rapid division of tumor cells (SFORCIN, 2016).

The cytotoxic activity of natural killer cells against murine lymphoma increased with the

Use of propolis for 3 days (SFORCIN, 2007). The presence of tumor suppressor proteins in the ester.

Phenethyl alcohol from caffeic acid causes apoptosis of C6 glioma cells (WATANABE et al., 2011; SFORCIN,

2016). Caffeic acid and esters, as well as diterpenoids and phenolic compounds have

destructive capacity against tumor cells. The antitumor effect of propolis is due to the function

combined with its polyphenolic constituents (SFORCIN, 2007). The decrease in the production of

Glutathione in the tumor cell, due to radiation, is consequently replenished by propolis, as

the synthesis of glutathione in hematopoietic tissues. Propolis originating from Türkiye acts as

antitumor activity, increasing tumor cell death, also showed a delay in leucine, thymidine and

uridine to become cancer-causing cells, restricting DNA synthesis (WATANABE)

et al., 2011).

3.4.10 Wound healing activity of propolis

The components of propolis also have therapeutic properties in tissue repair.

and regeneration of lesions (KUROPATNICKI et al., 2013). These are due to their effect

immunomodulatory, anti-inflammatory and antimicrobial characteristics (MARTINOTTI; RANZATO,

2015; SFORCIN, 2016). It was also noted that propolis decreases the amount of free radicals in the inflammatory

lesion and increases the development of collagen and its constituents (KRÓL et al., 2013;

MARTINOTTI; RANZATO, 2015). Accelerates different enzymatic reactions, metabolism of

cells, blood circulation, and also the formation of collagen fibers, due to the presence of

bioflavonoids, arginine, vitamin C, provitamin A, B complex, as well as some minerals

(PAROLIA et al., 2010).

4 CONCLUSION

Propolis is a natural product of the beehive derived from plant exudates. It has been

used as a traditional remedy for various ailments due to its biological activities and



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

pharmacological. Although, more than 350 constituents have been identified in samples of Propolis, but its biological activity is mainly due to a few substances (such as flavonoids, Propolis has a wide range of components including terpenes, caffeic acid, ferulic acid, CAPE, and coumaric acids and esters. multispectral properties such as antimicrobial activity against a wide range of microorganisms (bacteria, fungi and viruses), anti-inflammatory, anesthetic, healing, vasoprotective, Antioxidant, antitumor, anti-ulcer, hepatoprotective, and immunomodulatory.

Propolis is becoming increasingly popular due to its potential role in contribution to human health. Despite its beneficial activities, the most challenging problem This is due to uncertainty regarding its correct dosage and safety, as well as the chemical composition of propolis. It varies greatly due to differences in collection time, vegetation, and geographic location; therefore, the Biological activities of propolis collected in different phytogeographic areas and collection time. They also vary greatly, therefore defining the dosage is a difficult task. Furthermore, few individuals exhibit hypersensitivity to propolis (adverse effects such as rhinitis, conjunctivitis, rashes) skin conditions and bronchospasm), therefore, individuals who are allergic or hypersensitive to any of its Those involved should avoid using propolis or its supplements.

In conclusion, the development of new propolis compounds from propolis Originating from diverse geographical sources, this is vital in controlling various pathogenic diseases. A literature review does not exhaust the subject; on the contrary, it suggests that new and original research on the topic is needed. Propolis should be further explored for its potential properties against... human pathogens.

REFERENCES

AKCA, EA et al. *The Comparative Evaluation of the Antimicrobial Effect of Propolis with Chlorhexidine against Oral Pathogens: An In Vitro Study*. Biomed Research International, [SI], v. 2016, p. 1-8, Feb. 2016.

AL-ANI, I. et al. *Antimicrobial Activities of European Propolis Collected from Various Geographic Origins Alone and in Combination with Antibiotics*. Medicines, vol. 5, no. 1, p. 1-16, 2018.

AL-HARIRI, M. *Immune's-boosting agent: Immunomodulation potentials of propolis*. Journal of Family & Community Medicine, vol. 26, no. 1, p. 57-60, Jan./Apr. 2019.

ALMEIDA, EC; MENEZES, H. *Anti-inflammatory activity of propolis extracts: a review*. Journal of Venomous Animals and Toxins, vol. 8, no. 2, p. 13-21, May 2002.

ALYANE, M. et al. *Cardioprotective effects and mechanism of action of polyphenols extracted from propolis against doxorubicin toxicity*. Pakistan Journal of Pharmaceutical Sciences, vol. 21, no. 3, p. 201-209, Jul. 2008.

ANJUM, SI et al. *Composition and functional properties of propolis (bee glue): A review*. Saudi



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

Journal of Biological Sciences, vol. 26, no. 7, p. 1695-1703, nov. 2019.

BANSKOTA, AH et al. *Cytotoxic, hepatoprotective and free radical scavenging effects of propolis from Brazil, Peru, the Netherlands and China*. Journal of Ethnopharmacology, vol. 72, no. 1-2, p. 239-240, Sept. 2000.

BANSKOTA, AH et al. *Hepatoprotective and anti-Helicobacter pylori activities of constituents from Brazilian propolis*. Phytomedicine, vol. 8, no. 1, p. 16-23, Jan. 2001.

BARREIRAS, DG et al. *Efficacy of the antimicrobial action of jataí bee propolis extract (Tetragonisca angustula) against Gram-positive and Gram-negative bacteria*. Caderno de Ciências Agrárias, v. 12, p. 1-5, 2020.

BASTOS, EMAF; SANTANA, RA; CALÇA-COSTA, AGF; THIAGO, PS *Interaction between Apis mellifera L. and Baccharis dracunculifolia DC that favors green propolis production in Minas Gerais*. Brazilian Journal of Biology, v. 71, no. 3, p. 727-734, Aug. 2011.

BHAKUNI, GS et al. *Animal models of hepatotoxicity*. Inflammation Research, vol. 65, no. 1, p. 13-19, January 24, 2016.

BIESALSKI, HK et al. *Bioactive compounds: definition and assessment of activity*. Nutrition, vol. 25, no. 11, p. 1202–1205, Nov./Dec. 2009.

BREYER, HFE; BREYER, EDH; CELLA, I. *Production and processing of propolis*. Florianópolis: Epagri, 2016. 21 p.

BUENO-SILVA, B. et al. *The effect of seasons on Brazilian red propolis and its botanical source: chemical composition and bacterial activity*. Natural Product Research, vol. 31, no. 11, p. 1318-1324, jun. 2017.

CASTALDO, S.; CAPASSO, F. *Propolis, an old remedy used in modern medicine*. Phytotherapy, n. 73, suppl. 1, p. S1-S6, nov. 2002.

CHAN, GCF; CHEUNG, KW; SZE, DMY *The immunomodulatory and anticancer properties of propolis*. Clinical Reviews in Allergy & Immunology, vol. 44, no. 3, p. 262–273, jun. 2013.

CUESTA-RUBIO, O.; FRONTANA-URIBE, BA; RAMIREZ-APAN, T.; CARDENAS, J. *Polyisoprenylated benzophenones in Cuban propolis; biological activity of nemorosone*. Zeitschrift für Naturforschung, vol. 57, no. 3-4, p. 372-378, Mar. /abr. 2002.

DE LUCA, MP et al. *The anti-caries activity and toxicity of an experimental propolis-containing varnish*. Brazilian Oral Research, vol. 31, p. e45, mar. 2017.

DELAPLANE, KS *Honey Bees and Beekeeping*. Georgia (US): US Department of Agriculture, 2010.

DIDARAS, NA et al. *Antimicrobial Activity of Bee-Collected Pollen and Beebread: State of the Art and Future Perspectives*. Antibiotics, vol. 9, no. 11, p. 811, nov. 2020.

EL-GUENDOUIZ, S. et al. *Antioxidant and diuretic activity of co-administration of Capparis spinosa honey and propolis in comparison to furosemide*. Asian Pacific Journal of Tropical Medicine, vol. 10, no. 10, p. 974-980, Oct. 2017.



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

FERREIRA, JM et al. *New propolis type from north-east Brazil: chemical composition, antioxidant activity and botanical origin*. Journal of the Science of Food and Agriculture, vol. 97, no. 11, p. 3552-3558, Aug. 2017.

FISCHER, G.; HUBNER, SO; VARGAS, GD; VIDOR, T. *Immunomodulation by propolis*. Archives of the Biological Institute, v. 75, n. 2, p. 247-253, Apr./Jun. 2008.

FRANCHIN, M. et al. *Vestitol isolated from Brazilian red propolis inhibits neutrophils migration in the inflammatory process: elucidation of the mechanism of action*. Journal of Natural Products, vol. 79, no. 4, p. 954–960, apr. 2016.

GIMÉNEZ-CASSINA, B.; SCHMIDT, EM; EBERLIN, MN; SAWAYA, ACHF
Phytochemical markers of different types of red propolis. Food Chemistry, vol. 146, p. 174-180, Mar. 2014.

GÓMEZ-CARAVACA, A. et al. *Advances in the analysis of phenolic compounds in products derived from bees*. Journal of Pharmaceutical and Biomedical Analysis, vol. 41, no. 4, p. 1220–1234, jun. 2006.

HUANG, S. et al. *Recent advances in the chemical composition of propolis*. Molecules, vol. 19, no. 12, p. 19610–19632, nov. 2014.

HUANG, SS et al. *Antiarrhythmic effect of caffeic acid phenethyl ester (CAPE) on myocardial ischemia/reperfusion injury in rats*. Clinical Biochemistry, vol. 38, no. 10, Oct. 2005.

HUSSAIN, AI et al. *Seasonal variation in content, chemical composition and antimicrobial and cytotoxic activities of essential oils from four Mentha species*. Journal of the Science of Food and Agriculture, vol. 90, no. 11, p. 1827-1836, Aug. 2010.

KAISEROVÁ, H. et al. *Flavonoids as protectors against doxorubicin cardiotoxicity: Role of iron chelation, antioxidant activity and inhibition of carbonyl reductase*. Biochimica et Biophysica Acta – Molecular Basis of Disease, v. 1771, no. 9, p. 1065-1074, Sept. 2007.

KOMERICKI, P.; KRÄNKE, B. *Maculopapular exanthem from propolis: case report and review of systemic cutaneous and non-cutaneous reactions*. Contact Dermatitis, vol. 61, no. 6, p. 353–355, Dec. 2009.

KRÓL, W. et al. *Propolis: properties, application, and its potential*. Evidence-Based Complementary and Alternative Medicine, v. 2013, 807578, 2013.

KUROPATNICKI, AK; SZLISZKA, E.; KROL, W. *Historical Aspects of Propolis Research in Modern Times*. Evidence-Based Complementary and Alternative Medicine, v. 2013, Article ID 964149, 11 p., apr. 2013.

KÜÇÜK, M. et al. *Biological activities and chemical composition of three honeys of different types from Anatolia*. Food Chemistry, vol. 100, no. 2, p. 526–534, jun. 2007.

LABSKÁ, K.; PLODKOVÁ, H.; PUMANNOVÁ, M.; SENSCH, KH *Antiviral activity of propolis special extract GH 2002 against Varicella zoster virus in vitro*. Pharmazie, vol. 73, no. 12, p. 733-736, Dec. 2018.

LEWIS, CT; SHORT, C. *Oxford Latin Dictionary*. 2nd ed. Oxford (UK): Oxford University Press, 2012. 2368 p.



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

LOTFY, M. *Biological activity of bee propolis in health and disease*. Asian Pacific Journal of Cancer Prevention, vol. 7, no. 1, p. 22–31, Jan./Mar. 2006.

MACHADO, B. et al. *Propolis as an alternative in prevention and control of dental cavity*. Journal of Apitherapy, v. 1, no. 2, p. 47-50, Dec. 2016.

MARIN, F. et al. *Changes in nutraceutical composition of lemon juices according to different industrial extraction systems*. Food Chemistry, vol. 78, no. 3, p. 319–324, Aug. 2002.

MARTINOTTI, S.; RANZATO, E. *Propolis: a new frontier for wound healing?* Burns Trauma, vol. 3, no. 9, p. 1-12, Jul. 2015.

NICHITOI, MM et al. *Polyphenolics profile effects upon the antioxidant and antimicrobial activity of propolis extracts*. Scientific Reports, vol. 11, p. 20113, nov. 2021.

NOLKEMPER, S.; REICHLING, J.; SENSCH, KH; SCHNITZLER, P. *Mechanism of herpes simplex virus type 2 suppression by propolis extracts*. Phytomedicine, vol. 17, no. 2, p. 132-138, Feb. 2009.

ORSI, RO et al. *Susceptibility profile of Salmonella against the antibacterial activity of propolis produced in two regions of Brazil*. Journal of Venomous Animals and Toxins including Tropical Diseases, vol. 11, no. 2, p. 109-116, jun. 2005.

PARK, SG et al. *Evaluation of anti-allergic properties of caffeic acid phenethyl ester in a murine model of systemic anaphylaxis*. Toxicology and Applied Pharmacology, vol. 226, n. 1, p. 22-29, Jan. 2008.

PAROLIA, A.; THOMAS, MS; KUNDABALA, M.; MOHAN, M. *Propolis and its potential uses in oral health*. International Journal of Medicine and Medical Sciences, vol. 2, no. 7, p. 210-215, Jul. 2010.

PICCINELLI, AL et al. *Cuban and Brazilian red propolis: botanical origin and comparative analysis by HPLC-PDA/ ESI-MS/ MS*. Journal of Agricultural and Food Chemistry, vol. 59, no. 12, p. 6484-6491, jun. 2011.

POBIEGA, K. et al. *Comparison of the antimicrobial activity of propolis extracts obtained by means of various extraction methods*. Journal of Food Science and Technology, vol. 56, no. 12, p. 5386–5395, Dec. 2019.

POPOVA, MP et al. *Terpenes with antimicrobial activity from Cretan propolis*. Phytochemistry, vol. 70, no. 10, p. 1262-1271, jul. 2009.

RAO, PV; KRISHNAN, K. T.; SALEH, N.; GAN, S. *Biological and therapeutic effects of honey produced by honey bees and stingless bees: a comparative review*. Brazilian Journal of Pharmacognosy, v. 26, no. 5, p. 657–664, Sept./Oct. 2016.

REGUEIRA NETO, MS et al. *Seasonal variation of Brazilian red propolis: antibacterial activity, synergistic effect and phytochemical screening*. Food and Chemical Toxicology, vol. 107, p. 572-580, Sep. 2017.

SALATINO, A.; FERNANDES-SILVA, CC; RIGHI, AA; SALATINO, MLF *Propolis*



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

research and the chemistry of plant products. Natural Product Reports, vol. 28, no. 5, p. 925-936, May 2011.

SALOMÃO, K. et al. *Brazilian green propolis: effects in vitro and in vivo on Trypanosoma cruzi*. Evidence-Based Complementary and Alternative Medicine, v. 2011, n. 2011, 185918, Feb. 2011.

SCHNITZLER, P. et al. *Antiviral Activity and Mode of Action of Propolis Extracts and Selected Compounds*. Phytotherapy Research, vol. 24, suppl. 1, p. S20-S28, Jan. 2009.

SFORCIN, JM et al. *Seasonal effect on Brazilian propolis antibacterial activity*. Journal of Ethnopharmacology, vol. 73, no. 1-2, p. 243-249, nov. 2016.

SFORCIN, JM *Propolis and the immune system: a review*. Journal of Ethnopharmacology, vol. 113, no. 1, Aug. 2007.

SHARIFI-RAD, M. et al. *Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases*. Frontiers in Physiology, vol. 11, Jul. 2020

SHIMIZU, T. et al. *Anti-influenza virus activity of propolis in vitro and its efficacy against influenza infection in mice*. Antiviral Chemistry & Chemotherapy, vol. 19, no. 1, p. 7-13, Mar. 2008.

SIHERI, W.; ALENEZI, S.; TUSIIMIRE, J.; WATSON, DG *The Chemical and Biological Properties of Propolis*. In: ALVAREZ-SUAREZ, J. (ed.). *Bee Products – Chemical and Biological Properties*. New York: Springer, 2017.

SOUZA, EA et al. *Effects of seasonal variations and collection methods on the mineral composition of propolis from Apis mellifera Linnaeus beehives*. Brazilian Journal of Biology, v. 76, no. 2, p. 396-401, June 2016.

SOUZA, FBR; FISCHER, G.; VARGAS, GD. *Antimicrobial effect of propolis against infectious agents of veterinary interest*. Science and Animal Health, v. 1, n. 1, p. 24-37, Jul./Dec. 2013.

SUNG, SH et al. *External use of propolis for oral, skin, and genital diseases: a systematic review and meta-analysis*. Evidence-Based Complementary and Alternative Medicine, v. 2017, e8025752, Feb. 2017.

TORETI, VC et al. *Recent Progress of Propolis for Its Biological and Chemical Compositions and Its Botanical Origin*. Evidence-Based Complementary and Alternative Medicine, v. 2013, e697390, apr. 2013.

VALENCIA, D. et al. *Seasonal effects on chemical composition and biological activities of Sonoran propolis*. Food Chemistry, vol. 131, no. 2, p. 645-651, Mar. 2012.

VECCHI, E.; DRAGO, L. *Propolis' antimicrobial activity: what's new?* Le Infezioni in Medicina, v. 15, no. 1, p. 7-15, Mar. 2007.

VEIGA, RS et al. *Artepillin C and phenolic compounds responsible for antimicrobial and antioxidant activity of green propolis and Baccharis dracunculifolia DC*. Journal of Applied Microbiology, vol. 122, no. 4, p. 911-920, apr. 2017.

VOLPI, N. *Separation of flavonoids and phenolic acids from propolis by capillary zone electrophoresis*. Electrophoresis, vol. 25, no. 12, p. 1872–1878, jun. 2004.



Year V, v.2 2025 | Submission: November 14, 2025 | Accepted: November 16, 2025 | Publication: November 18, 2025

WAGH, VD *Propolis: A Wonder Bees Product and Its Pharmacological Potentials*. Advances in Pharmacological and Pharmaceutical Sciences, vol. 2013, e308249, Dec. 2013.

WANG, HX et al. *Immune mechanisms of Concanavalin A model of autoimmune hepatitis*. World Journal of Gastroenterology, vol. 18, no. 2, p. 119-125, Jan. 2012.

WATANABE, MAE et al. *Cytotoxic constituents of propolis inducing anticancer effects: a review*. The Journal of Pharmacy and Pharmacology, vol. 63, n. 11, p. 1378–1386, nov. 2011.

WILLIAMS, CD et al. *Mouse strain-dependent caspase activation during acetaminophen hepatotoxicity does not result in apoptosis or modulation of inflammation*. Toxicology and Applied Pharmacology, vol. 257, n. 3, p. 449-458, Dec. 2011.

WOO, KJ et al. *Chrysin suppresses lipopolysaccharide-induced cyclooxygenase-2 expression through the inhibition of nuclear factor for IL-6 (NF-IL6) DNA-binding activity*. FEBS Letters, vol. 579, no. 3, p. 705-711, Jan. 2005.

XUAN, H. et al. *Anti-inflammatory effects of Chinese propolis in lipopolysaccharide-stimulated human umbilical vein endothelial cells by suppressing autophagy and MAPK/NF- κ B signaling pathway*. Inflammopharmacology, v. 27, p. 561-571, Jan. 2019.

YE, M. et al. *Alterations in the Transcriptional Profile of the Liver Tissue and the Therapeutic Effects of Propolis Extracts in Alcohol-induced Steatosis in Rats*. Annals of the Brazilian Academy of Sciences, v. 91, no. 3, e20180646, Aug. 2019.

YINKANG, L. et al. *Galangin and Pinocembrin from Propolis Ameliorate Insulin Resistance in HepG2 Cells via Regulating Akt/mTOR Signaling*. Evidence-Based Complementary and Alternative Medicine, v. 2018, p. 7971842, Oct. 2018.

YOSEFI, S. et al. *Chrysin-Enhanced Cytotoxicity of 5-Fluorouracil-Based Chemotherapy for Colorectal Cancer in Mice: Investigating its Effects on Cyclooxygenase-2 Expression*. Brazilian Journal of Pharmaceutical Sciences, v. 58, e19381, 2022.

ZABAIU, N. et al. *biological properties of propolis extracts: something new from an ancient product*. Chemistry and Physics of Lipids, vol. 207, p. 214-222, Oct. 2017.