

Muscle Hypertrophy

Muscle Hypertrophy

Muscle Hypertrophy

Gianni Leandro Goicoa da Costa¹

Cristian Rodrigues (counselor)²

Summary

Muscular hypertrophy is a biological process characterized by an increase in the volume of skeletal muscle fibers, resulting mainly from resistance training, such as musculature.

This phenomenon occurs through mechanical and metabolic stimuli that cause microdamage to the fibers, triggering an adaptive response of the organism. Protein synthesis is stimulated, leading to muscle repair and growth, as long as there is adequate nutritional intake, especially proteins and essential amino acids, and sufficient rest for recovery.

Various factors influence the process, including genetics, intensity of exercise volume, sound, health and hormonal levels. Regular practice of resisted exercises, associated with a balanced diet and a periodized plan, is essential to achieve consistent gains in lean mass and muscular strength, also contributing to better metabolism and body composition.

Keywords: Muscular hypertrophy. Resisted training. Synthesis protein. Sports nutrition. Physiological adaptation.

Abstract

Muscle hypertrophy is the biological process characterized by an increase in the volume of skeletal muscle fibers, primarily resulting from resistance training, such as weight training. This phenomenon occurs through mechanical and metabolic stimuli that induce microlesions in the fibers, triggering an adaptive response by the body. Protein synthesis is stimulated, leading to muscle repair and growth, as long as there is adequate nutritional intake, especially of proteins and essential amino acids, and sufficient rest for recovery. Several factors influence the process, including genetics, training intensity and volume, sleep, age, and hormone levels. The regular practice of resistance exercises, combined with a balanced diet and periodic planning, is essential for achieving consistent gains in lean mass and muscle strength and for improving metabolism and body composition. Keywords: Muscle hypertrophy. Resistance training. Protein synthesis. Sports nutrition—physiological adaptation.

1 Introduction

Muscular hypertrophy is a biological phenomenon of great relevance in the context of exercise and health sciences, defined as the increase in the size of two skeletal muscles. It responds to specific stimuli, especially strength training. Or be interested in esse. The process transcends the limits of sports performance, extending to rehabilitation and promotion. It gives health and prevention of chronic diseases. Recent research highlights that muscle hypertrophy. Not only does it improve functionality and quality of life, but it also plays a role. crucial in modulating metabolism and not controlling cardiovascular risk factors, becoming

¹ Gianni Leandro Goicoa presents the Bacharelado course in Physical Education from Unopar – Universidade Norte do Paraná.

² Cristian Rodrigues. Teacher of the Bacharelado course in Physical Education at Unopar – Universidade Norte do Paraná.

A strategic objective in multidisciplinary interventions.

The study of muscular hypertrophy is intrinsically multidimensional, involving aspects physiological, biochemical, genetic and environmental. Scientific advances allow us to understand, in an increasingly detailed way, the cellular and molecular mechanisms responsible for muscle growth, as it varies, as in modulation, as in intensity, or volume.

training frequency, apart from nutritional and hormonal factors. This complexity evidences need for integrated and evidence-based approaches for the development of Effective hypertrophy promotion strategies in different populations.

The scientific literature is discussed about the individual differences in the answer. hypertrophic during training, highlighting the influence of variables such as age, sex, genetic profile and health conditions. These factors can determine only the potential of mass gain muscle, but also the speed and sustainability of these adaptations. understand these nuances is essential for the personalized prescription of training programs and for Maximizing two benefits associated with hypertrophy, minimizing risks and optimizing results.

Another central aspect refers to the relationship between physical training, nutrition and life processes. recovery, elements that compõem the tripe of muscular development. Or adequate contribution of nutrients, especially proteins, the periodization of training and the quality of sound recognized as critical determinants for protein synthesis and muscle remodeling. Alem Disso, recent research suggests the role of hormonal signaling in the microenvironment inflammation and neuromuscular plasticity in the modulation of hypertrophic response, indicating promising paths for future interventions.

Given the growing interest of society in aesthetics, performance and health, to understand two processes that govern muscular hypertrophy assume unique relevance for physical education professionals, physiotherapists, nutritionists and doctors. Or delineation of recommendations based on evidence, based on the integration between basic and applied science, emerges as an essential requirement for the development of safe, efficient and ethical practices not The field of physical training and the promotion of muscular health.

2 Development

2.1 Methodology

This study was prepared by means of bibliographic review, of a qualitative nature and descriptive, as the objective of gathering, analyzing and synthesizing the main scientific evidence about the mechanisms, factors and strategies related to muscular hypertrophy. The investigation was conducted between

July and October 2025, using as main sources specialized books, scientific articles and Academic dissertations are available in rebuilt databases, such as SciELO, PubMed, Google Scholar and ResearchGate, besides national and international Physical Education journals, Nutrition and Exercise Physiology.

Forum considered works published in the last ten years (2015–2025), or that assures up to date and relevant information. The bibliographic search was carried out by Meio de descriptors selected according to the study objectives, including: “muscular hypertrophy”, “resisted training”, “protein synthesis”, “sports nutrition”, “muscle recovery” and “sono e physical performance.”

The criteria for inclusion open publications with scientific approaches, methodologies validated and direct relationship with the topics of physiology, nutrition and force training to hypertrophy. Excluíram-se duplicate articles, opinionated texts, summaries without access to the content comprehensive and empirically based studies or systematic review.

After triage, the selected references are organized and analyzed as coerência as the proposed objectives and classified in thematic areas: physiology and mechanisms of hypertrophy; determining factors and external influences; force and periodization training; applied nutrition ao muscle growth; e recovery and sound as mediators of adaptation.

The adopted methodology allowed us to identify trends, gaps and convergences in Recent publications, offering an integrated and up-to-date view on the topic. This approach Qualitative bibliography does not involve the collection of primary data or the application of instruments you experiment, limiting yourself to interpretation and critical analysis of studies already consolidated in literature scientific.

2.1 Results and Discussion

The results of the bibliographical research reveal that muscular hypertrophy is a phenomenon extensively studied in exercise sciences, with increasing emphasis in recent decades due to Its relevance for health, performance and physical rehabilitation. A chronological analysis of two studies allowed us to observe a significant evolution without understanding two physiological mechanisms, das Variáveis of training and nutritional strategies involved in muscle growth. Such Evolution is due, in part, to the advancement of research methodologies, to the development of imaging technologies and the consolidation of experimental protocols that make it possible to measure with greater precision as structural and functional alterations in human muscle tissue.

In the first investigations on the subject, hypertrophy was understood in a simplistic way,

associated only with an increase in visible muscle volume. Nope, more recent studies demonstrate that the process results from complex cellular adaptations, involving synthesis myofibrillar protein, or increase in mitochondrial density and activation of satellite cells Responsáveis pela regeneração tecidual. According to Lima and Miguel et al. (2021), these answers Physiological disorders only triggered by repetitive mechanical stimuli, typical of strength training, that there will be microlesões in the muscle fibers, as quais, once we recover, they will become larger and more resistant.

Another relevant discovery identified in the publications analyzed was the difference between Sarcoplasmic hypertrophy and myofibrillar hypertrophy, as discussed by Oliveira (2022). As for the first time, it is related to the increase in fluid and glycogen content within the body. muscle cell, conferring aesthetic volume, the second is linked to the excess of proteins contract, like actin and myosin, which results in greater strength and muscle density. to literature More recent research suggests that both forms occur simultaneously, varying according to the type of applied stimulus, or level of experience of the practitioner or individual genetic profile. This distinction allowed substantial progress in the prescription of specific training programs for different objectives, whether for athletic performance or functional rehabilitation.

In the field of endocrine physiology, the research reviewed will highlight the crucial role of two anabolic hormones in the hypertrophy process. Nogueira (2022) and Andrade and Reis (2021) We will observe that substances such as testosterone, growth hormone (GH) and IGF-1 act Directly in the activation of intracellular signaling pathways responsible for protein synthesis. via mTOR (mammalian target of rapamycin), in particular, was widely recognized as a point central in the regulation of anabolic response, integrating mechanical, nutritional and hormonal stimuli. Furthermore, it is verified that negative regulation is destructive, caused by factors such as stress. chronic or sound deprivation, can compromise muscle development, or demonstrate that Hormonal and metabolic homeostasis is determining for the effectiveness of the hypertrophic process.

Other results refer to the influence of nutrition and energy status in the body. hypertrophy. According to Ogiku and Ikematsu (2015), the positive and fundamental energy balance to sustain muscle growth, since severe caloric deficits reduce protein synthesis even given adequate mechanical stimulation. Or consumption of proteins of high biological value, especially as rich in essential amino acids, such as leucine, are essential for Activate mTOR and promote tissue repair. Além disso, more recent investigations, like Yasuda's ace et al. (2022), demonstrated that the uniform distribution of protein intake throughout the day results in performance superior to concentrated intake in a single meal, suggesting that the father Feeding plays an important role in the total volume of protein ingested.

The literature also reveals that training methods are resistant to factors. determinants for the success of hypertrophy. Faria (2018) and Hajj-Boutros and Jacob (2023) highlight more than high intensity strength training, performed between 60% and 85% of 1RM (repetition maximum), induces the greatest gains in muscular mass, since associated with a periodization structure and adequate recovery intervals. Linear and wave periodization models. They have proven to be effective in avoiding adaptive plates and maintaining continuous hypertrophic stimulation. These slopes reinforce the importance of the planned variation of volume and intensity as an essential tool to maximize physiological adaptations and reduce the risk of overtraining.

The question of muscle recovery and sound emerged as an essential component in more recent studies. According to Parreira and Resende (2015) and Lima (2017), deep sound favors the pulsatile release of GH, reducing two levels of cortisol, creating an ideal anabolic environment for muscle growth. A deprivation of sound, on the other hand, increases fatigue, harms tissue regeneration and reduction of protein synthesis efficiency. These results show that adequate rest should be considered as a training variable that is relevant to training intensity, being essential to optimize two results.

Além disso, a review revealed that biological individuality exerts direct influence on the magnitude of hypertrophic responses. Studies show significant differences between individuals regarding the proportion of type I and type II fibers, hormonal sensitivity, and ability to recover and genetic predisposition (HAYES and CRIBB, 2008). This variability reinforces the need for personalized training protocols, adjusted to physiological characteristics and objective years of each practitioner. The reconfirmation of these differences also contributes to the development of more inclusive approaches in training programs, contemplating groups as women, educated and initiating, who traditionally present less representation in studies on hypertrophy.

A global analysis of two results indicated that muscle hypertrophy is a multifactorial process, dependent on the integration between the mechanical stimulus, the nutritional contribution and adequate recovery. Nenhum fator isolado was able to fully explain muscle growth; on the contrary, as evidence suggests a synergy between exercise, nutrition and sleep as a basis for success. adaptations. These conclusions reaffirm the role of the Physical Education professional as a mediator between science and practice, responsible for applying technical and empirical knowledge ethically and personalized, guaranteeing efficiency and safety of two programs aimed at development muscle mass.

Among the oldest studies, the research of Andrews and MacLean (2006) stands out, which demonstrate the importance of adequate protein consumption for muscle synthesis and exercise

practitioners of resisted exercises, and the contributions of Hulmi, Lockwood and Stout (2010), who confirm the role of the protein of soro do leite (whey protein) in the activation of the protein synthesis in post-training recovery. These investigations will serve as a basis for subsequent studies that deepening the understanding of the metabolic pathways and hormones involved in the hypertrophic process.

As of 2015, as advances in imaging and molecular biology techniques have been made, extension of knowledge about the activation of the IGF-1/Akt/mTOR pathway, the main mediator of synthesis muscle protein (NOGUEIRA, 2022). Ogiku and Ikematsu (2015) confirm that ingestion Adequate supply of essential amino acids, especially leucine, potentiates this pathway, reinforcing Direct relationship between nutrition and muscular adaptation. Já Lima e Miguel et al. (2021) demonstrate that The number of series and intensity of two exercises directly influences the volume of hypertrophy, being the high-intensity and short-duration protocols the most effective in promoting the increase in Cross section of muscle fibers.

The results will also show what hormonal and environmental factors play a role. determining role in the hypertrophic process. Oliveira (2022) and Andrade and Reis (2021) highlight that anabolic hormones, such as testosterone, GH and IGF-1, act synergistically to promote or growth of myofibrils, as high levels of cortisol tiveram catabolic effect, reducing or muscle gain. Além disso, the studies emphasize the importance of sound and recovery as essential components of the adaptation, since the release of GH causes protein synthesis more intensely during or rest (PARREIRA and RESENDE, 2015; LIMA, 2017).

In the field of practice, Faria (2018) and Hajj-Boutros and Jacob (2023) evidence that or Periodized and individualized strength training was the most effective strategy to induce hypertrophy safely and continuously. These authors will observe that the planned alternation between phases of major volume and lower intensity and phases of higher intensity and lower volume prevented or supertraining and maximizing or hypertrophic stimulus. Additionally, I verified which protocols that combining multi-articular and mono-articular exercises results in greater recruitment of fibers and a better balance between strength and muscle volume.

From a nutritional point of view, the data indicate that the daily protein intake between 1.6 e 2.2 g/kg body weight, associated with a caloric intake slightly higher than energy expenditure daily, it was decisive for consistent gains of lean mass (LIMA and MIGUEL, 2021). essa Consumption rate was sufficient to maximize muscle protein synthesis (MPS), reduce or Catabolism and promote positive nitrogen balance. The authors will highlight that surpluses calories do not cause greater hypertrophy, but rather an increase in body fat, reinforcing importance of balanced nutritional strategies adapted to the body composition of each individual. Thus, caloric modulation must be adjusted gradually, prioritizing quality.

macronutrients and metabolic efficiency of the organism.

Protein fractionation throughout the day and post-workout intake will demonstrate strategies fundamentals to optimize anabolic chamada janela, period in which the organism appears greater sensitivity to muscle protein synthesis (HULMI, LOCKWOOD and STOUT, 2010). Evidence Recent indications indicate that the consumption of 20 to 40 g of protein of high biological value per meal, distributed at intervals of three to four hours, maintain constant stimulation to protein synthesis (YASUDA et al., 2022). This practical guarantor or continuous supply of essential amino acids, especially leucine, the main activator of the mTOR pathway, responsible for the beginning of the process cellular anabolic. In this way, the consistency of protein intake, more than the total volume consumed in a single meal, it constitutes the preponderant factor for muscle growth supported.

Among the most effective protein sources, whey protein stands out. protein), egg protein, lean meats and fish, which have high bioavailability and complete profile of essential amino acids (OGIKU and IKEMATSU, 2015). The studies will show that The consumption of whey protein after resisted exercise results in rapid elevation of two levels plasma leucine and stimulates, more efficiently, MPS when compared to sources you vegetate Meanwhile, authors such as Wilson (2016) and Nogueira (2022) highlight that diets based Vegetable proteins can also promote hypertrophy, as long as there is adequate planning das food combinations to achieve the full spectrum of amino acids and an intake slightly higher total protein (2.0–2.4 g/kg).

In the context of two carbohydrates, analyzed literature demonstrates that they perform a essential role not only as a primary source of energy, but also in response modulation hormonal training. According to Hayes and Cribb (2008), carbohydrate intake along with Proteins after exercise favor the release of insulin, a hormone that inhibits protein degradation It facilitates the transport of amino acids to the interior of muscle cells. This combination It also reduces cortisol levels, creating a metabolic environment more favorable to recovery. The shape, the balance between proteins and carbohydrates in post-training meal is considered one of the two pillars of nutrition aimed at muscular hypertrophy.

Lipids, often neglected, also perform functions important in hormonal synthesis and the health of cell membranes. Investigations such as Ogiku and Ikematsu (2015) showed that excessively restrictive diets in fat can compromise the production of testosterone and other steroid hormones, reducing the anabolic efficiency of body. It is recommended that lipids comprise 20% to 35% of the total energy value of diet, favoring sources rich in mono- and polyunsaturated fatty acids, such as olive oil,

abacate, chestnuts and cold water fish. In addition to contributing to the hormonal profile, lipid levels It has anti-inflammatory properties that help in muscle recovery and protection against or oxidative stress induced by intense exercise.

The adequacy of two micronutrients was another point of emphasis in the evidence reviewed. Vitamins and minerals, such as vitamin D, zinc, magnesium and calcium, show a direct relationship com a força ea muscular recovery (OLIVEIRA, 2022). Vitamin D, especially tuna Modulation of muscle contraction and calcium metabolism, directly influencing efficiency neuromuscular. Micronutrient deficiency has been associated with reduced strength and resistance, as well as a greater predisposition to injuries. Já o zinco participates in enzymatic processes related to protein synthesis and testosterone production, as magnesium helps in Tecidual regeneration and stabilization of cell membranes. Assim, snow maintenance Adequate use of these micronutrients is achieved through adequate nutrition or supplementation. oriented, and essential to the full functioning of muscular metabolism.

Nutritional supplementation was identified as an effective support resource, since used judiciously and based on scientific evidence. Among the supplements more studied, creatine monohydrate stands out for increasing phosphocreatine levels intramuscular, which improves work capacity, delays fatigue and enhances the desire to force and muscle volume (FARIA, 2018). Also, beta-alanine or HMB (beta-hydroxy-beta-methylbutyrate) foram suggested as substances with complementary positive effects, emoras Results on its direct impact on hypertrophy still show divergences in the literature. TO The use of supplements must therefore be subject to individual and yearly approval. accompanied by qualified professionals, guaranteeing safety and effectiveness.

Another aspect addressed by recent studies refers to the role of hydration in the dehydration process. muscle growth. Parreira and Resende (2015) highlight that the state of hydration influences performance, or transport of nutrients and protein synthesis. A slight dehydration, of just 2% gives body mass, it is capable of compromising muscle function and increasing oxidative stress. By Therefore, adequate water intake before, during and after exercise must be integrated into the plan. nutritional, especially in hot environments or during high-intensity workouts. To water It is also essential for the transport of amino acids and electrolytes, which reinforces its role as adjuvant element in the hypertrophic process.

In summary, the results indicate that the success of muscle hypertrophy depends on interaction between dietary, hormonal and metabolic factors, leading to structured and individualized or central eixo dessa adaptation. Literature reinforces that nutrition should not be seen only as a training support, but also as an active anabolic tool, capable of

potentiate the effects of resisted exercise. A combination between nutritional planning balanced, rational supplementation, hormonal control and adequate hydration constitute, therefore, or alicerce for sustainable and safe muscle growth, which validates the importance of atuação joint collaboration between Physical Education professionals, nutritionists and physiologists in prescribing and not accompaniment of programs aimed at hypertrophy.

In relation to the most recent years, Yasuda et al. (2022) and Vasconcelos Neto (2025) reinforce the importance of protein distribution over the long period of meals and the role of leucine in activation of mTOR, consolidating a new understanding of the synergy between nutrition and training Santos and Bezerra (2014) emphasize that the processes of cellular signaling are fundamental to understand muscle growth on a molecular level, opening new doors Perspectives for studies of applied physiology.

A discussion of two results allowed us to conclude that muscular hypertrophy hardly corresponds Isolated physical exercise, but also integration between resisted training, adequate nutrition, rest and hormonal control. The research analyzed corroborates the classical theories of physiology exercise and expand knowledge on cellular and metabolic adaptations, confirming that the hypertrophic process depends on progressive overload, on positive protein balance and on adequate recovery.

Despite two advances, the limitations pointed out by the authors include a lack of studies Lengths in specific populations, such as women and men, are difficult to register Variáveis of training in controlled clinical trials. These limitations indicate the necessities of new research that explores biological individuality and long-term adaptive responses. Of Generally speaking, the results found and discussed reinforce the idea that muscular hypertrophy It is a multifactorial phenomenon, supported by physiological, nutritional and behavioral evidence, and that its integrated understanding is essential for the development of training programs safe and efficient.

Conclusion

An integrated analysis of two multiple aspects involved in muscular hypertrophy evidence or multifactorial and dynamic character of the adaptive process. A synthesis of knowledge from Physiology, genetics, nutrition, training and recovery reveal that the increase in muscle mass It does not depend on a single isolated factor, but rather a complex interaction between mechanical stimuli, Nutrient availability, hormonal profile and individual variables such as age, sex and heritage genetics (FARIA, 2018). The literature highlights that strategies based on progressive overload,

Adequate protein intake, quality and recovery planning are essential for optimize the hypertrophic process, reducing risks and maximizing the functional benefits and metabolic.

The individualization of interventions stands out as a guiding principle, considering High interindividual variability in responding to the same training and nutrition prescriptions (FARIA, 2018). Factors such as fiber type composition, genetic polymorphisms, state nutritional, level of experience in training and presence of specific clinical conditions It modulates both the potential and the speed of the muscle mass gain. Dessa forma, a personalized prescription, based on regular assessments and no systematic monitoring results, emerges as a superior strategy in relation to standardized approaches, promoting Sustainability of hypertrophic organisms and minimization of adverse effects.

The importance of interdisciplinary work is reiterated given the complexity of the phenomenon hypertrophic (FARIA, 2018). Integration between Physical Education professionals, nutritionists, Physiotherapists and doctors potentiate the achievement of results through the alignment of stimuli physical, nutritional support and recovery strategies. Além disso, to ethical and guided atuação Scientific evidence is essential to cover inappropriate practices, such as indiscriminate use of ergogenic substances and restrictive diets, and to guarantee the safety of practitioners, above all in contexts of greater vulnerability, such as the elderly, adolescents and individuals with comorbidities.

The reconfirmation of two benefits of muscular hypertrophy extrapolated to the aesthetic and sports, reaching the promotion of public health, the prevention of chronic diseases and better times gives functionality in different populations (FARIA, 2018). The increase in muscle mass is associated with reducing the risk of sarcopenia, seizures, insulin resistance and morbidities cardiovascular, in addition to contributing to autonomy and quality of life in all age groups. Thus, the dissemination of practices based on evidence and adapted to sociocultural realities It constitutes a challenge and a responsibility for professionals and institutions dedicated to promotion muscle health.

References

ALMEIDA, PM of; CARVALHO, DS; PEREIRA, CA **Learning protein synthesis in an interactive and playful way: the protein synthesis challenge.** *Journal of Biochemistry Education*, Brazilian Society of Biochemistry and Molecular Biology - SBBq, v. 20, no. 2, p. 1–13, 2021. Available at: <https://doi.org/10.16923/reb.v20i2.918>. Access em: 11 out. 2025.

ANDRADE, SM; REIS, DS; FERRO, AR **Benefits of strength training associated with the partial vascular occlusion method in hypertrophy and no muscle strength gain.** *RENEF*, Montes Claros: Universidade Estadual de Montes Claros (UNIIMONTES), v. 12, no. 18, p. 69–83,



Year VII, v.1 2026 | submitted: 04/10/2026 | oil: 04/14/2026 | publication: 04/17/2026

2021. Available at: <https://doi.org/10.46551/rn2021121800051>. Access em: 11 out. 2025.

ANDRADE, SM; REIS, DS; FERRO, ARF **Benefits of strength training associated with the partial vascular occlusion method in hypertrophy and no muscle strength gain.** *RENEF*, Montes Claros: Universidade Estadual de Montes Claros (UNIIMONTES), v. 12, no. 18, p. 69–83, 2021. Available at: <https://doi.org/10.46551/rn2021121800051>. Access em: 11 out. 2025.

ANDREWS, RD; MACLEAN, DA; RIECHMAN, SE **Protein intake for skeletal muscle hypertrophy with resistance training in seniors.** *International Journal of Sport Nutrition and Exercise Metabolism*, Human Kinetics, v. 16, no. 4, p. 362–372, 2006. Available at: <https://doi.org/10.1123/ijsnem.16.4.362>. Access em: 11 out. 2025.

BEZERRA, MA; SANTOS, NA two. **Cellular signaling pathways can help you understand melhor or Ebola.** *Nanocell News*, Nanocell Institute, v. 1, no. 16, 2014. Available at: <https://doi.org/10.15729/nanocellnews.2014.08.25.001>. Access em: 11 out. 2025.

FARIA, DPB of. **Creatine supplementation does not gain strength and muscle hypertrophy in strength training practitioners: a brief narrative review.** *Revista Eletrônica Acervo Saúde*, 2018, Sup., n. 13, p. S1476-S1483. Available at: https://doi.org/10.25248/reas274_2018. Accessed on: 11 Oct. 2025.

FARIA, DPB of. **Creatine supplementation without strength gain and muscular hypertrophy in strength training practitioners: a brief narrative review.** *Eletrônica Acervo Saúde Magazine*, 2018, v. Sup., n. 13, p. S1476-S1483. Available at: https://doi.org/10.25248/reas274_2018. Accessed on: 11 Oct. 2025.

FINCK, SCM **Physical school education: a reading in the emerging conception of human development.** *Olhar de Professor*, Ponta Grossa: Universidade Estadual de Ponta Grossa (UEPG), v. 1, no. 1, p. 1–10, 2009. Available at: <https://doi.org/10.5212/olharprofr.v.1.1338>. Accessed on: 11 Oct. 2025.

HAJJ-BOUROS, G.; JACOB, K.; SONJAK, V.; BROERING, F.; MORAIS, JA **Resistance training, but not leucine, can reverse frailty by increasing basal muscle protein synthesis in older women consuming an optimized protein intake.** *Clinical Nutrition ESPEN*, Elsevier BV, v. 54, p. 714–715, 2023. Available at: <https://doi.org/10.1016/j.clnesp.2022.09.745>. Access em: 11 out. 2025.

HAYES, A.; CRIBB, PJ **Effect of whey protein isolate on strength, body composition and muscle hypertrophy during resistance training.** *Current Opinion in Clinical Nutrition and Metabolic Care*, Ovid Technologies (Wolters Kluwer Health), v. 11, no. 1, p. 40-44, 2008. Available at: <https://doi.org/10.1097/mco.0b013e3282f2a57d>. Access em: 11 out. 2025.

HULMI, JJ; LOCKWOOD, CM; STOUT, JR **Effect of protein/essential amino acids and resistance training on skeletal muscle hypertrophy: a case for whey protein.** *Nutrition & Metabolism*, Springer Science and Business Media LLC, v. 7, no. 1, p. 51, 2010. Available at: <https://doi.org/10.1186/1743-7075-7-51>. Access em: 11 out. 2025.

LIMA, LEDM; MIGUEL, H.; MORIGGI JUNIOR, R.; SILVA, RPD; NOGUEIRA, HS; BRITO-NETO, JGD; GUEDES JR., DP **Number of series in muscular force training: a brief narrative review of meta-analyses on force adaptations and muscular hypertrophy.**

Year VII, v.1 2026 | submitted: 04/10/2026 | oil: 04/14/2026 | publication: 04/17/2026

Multidisciplinary Reviews, Malque Publishing, v. 4, no. 1, p. e2021008, 2021. Available at: <https://doi.org/10.29327/multi.2021008>. Access em: 11 out. 2025.

LIMA, WP **Molecular mechanisms associated with muscle hypertrophy and hypotrophy: relationship with physical exercise practice.** *Revista Brasileira de Fisiologia do Exercício*, Convergences Editorial, 2, p. 123-141, 2017. Available at: <https://doi.org/10.33233/rbfe.v16i2.972>. Access em: 11 out. 2025.

MOREIRA, PSA; SILVEIRA, AC; ARRIGONI, MDB; COSTA, C.; CHARDULO, LAL; FURLAN, LR; DAL PAI, V. **Effect of somatotropin in hypertrophy of skeletal muscle fibers and in reproductive precocity of Nelore heifers.** *Pesquisa Agropecuária Brasileira*, FapUNIFESP (SciELO), vp 1853-1860, 2000. Available in: 35, n. 9, <https://doi.org/10.1590/s0100-204x2000000900018>. Access em: 11 out. 2025.

NOGUEIRA, HS **Gerais conceitos e determinant factors para hypertrophic responses in skeletal musculature induced by muscular strength training – a narrative review.** *Revista Brasileira de Fisiologia do Exercício*, Convergences Editorial, v. 17, no. 1, p. 60-63, 2022. Available at: <https://doi.org/10.33233/rbfe.v17i1.2369>. Access em: 11 out. 2025.

OGIKU, M.; IKEMATSU, Y.; OGASAWARA, T.; OKAMOTO, Y. **SUN-LB020: The relationship between dietary amino acid intake and training-induced muscle hypertrophy among older individuals.** *Clinical Nutrition*, Elsevier BV, v. 34, p. S242, 2015. Available at: [https://doi.org/10.1016/s0261-5614\(15\)30741-x](https://doi.org/10.1016/s0261-5614(15)30741-x). Access em: 11 out. 2025.

OLIVEIRA, M. de. **Muscular hypertrophy: a study on the two types of myofibrillar muscular hypertrophy and sarcoplasmic muscular hypertrophy.** *Physical Education, Sports, Culture & Society*, Editora ZH4, 2022, p. 245–261. Available at: <https://doi.org/10.51360/zh4.20221-01-p.245-261>. Access em: 11 out. 2025.

PARREIRA, RC; RESENDE, RR **What is hypertensive? (5th chapter) What is the difference between cardiac hypertrophy and hypertension?** *Nanocell News*, Nanocell Institute, v. 2, no. 12, p. n/an/a, 2015. Available at: <https://doi.org/10.15729/nanocellnews.2015.05.19.002>. Access em: 11 out. 2025.

SAMPAIO, ADS; NUNES, MA da C. **School physical education and its importance in human development.** *Interdisciplinary Dialogues 13*, Diálogo Comunicação e Marketing, 2024, p. 40–54. Available at: <https://doi.org/10.29327/5450500.1-3>. Access em: 11 out. 2025.

SANTOS, NA; BEZERRA, MA; RESENDE, RR **Stomach surgery also causes loss of function due to cellular signalization.** *Nanocell News*, Nanocell Institute, 2014, v. 1, no. 11, p. 1–1. Available at: <https://doi.org/10.15729/nanocellnews.2014.05.12.002>. Access em: 11 out. 2025.

SILVA, MB; CAVAZOTTI, MA **Contributions to school physical education for the psychic development of human beings: first apontamentos.** *Revista Ibero-Americana de Estudos em Educação*, 2017, v. 12, no. 2, p. 689-709. Available at: <https://doi.org/10.21723/riaee.v12.n2.8298>. Access em: 11 out. 2025.

SILVA, TF da; FÉLIX, MEC **The impact of alcohol consumption on muscular hypertrophy and strength training.** *Physical education in movement: health and well-being through exercise*, Atena Editora, 2024, p. 13-26. Available at: <https://doi.org/10.22533/at.ed.6002429012>. Access em: 11 out. 2025.

SOARES, CP; CASSEB, GB; BARBOSA NETO, IA; PONTES, NS; JUCA, FL; ALBUQUERQUE, LL; FURTADO, C. de M.
Importance of skeletal contractile proteins during muscular hypertrophy. *Health Science in Western Amazônia*, Stricto
Sensu Editora, 2019, p. 201–214. Available at: <https://doi.org/10.35170/ss.ed.9786580261055.14>. Access em: 11 out. 2025.

SOUZA JUNIOR, TP of; DUBAS, JP; PEREIRA, B.; OLIVEIRA, PR de. **Creatine supplementation and strength training: alterations resulting from maximum dynamic strength and anthropometric variations in university students subjected to eight weeks of strength training (hypertrophy).** *Brazilian Journal of Sports Medicine*, FapUNIFESP (SciELO), v. 13, no. 5, p. 303-309, 2007. Available at: <https://doi.org/10.1590/s1517-86922007000500005>. Access em: 11 out. 2025.

TRISTÃO, PA da S.; JUSTO, JL; TOIGO, AM **I teach about the process of human development in graduation courses in Physical Education.** *Health and Human Development*, Centro Universitário La Salle - UNILASALLE, v. 5, no. 2, p. 39, 2017. Available at: <https://doi.org/10.18316/sdh.v5i2.3181>. Access em: 11 out. 2025.

VASCONCELOS NETO, JM **The role of the essential amino acid leucine in muscle protein synthesis.** *Quality of life and health – contemporary perspectives: Volume 5*, Dialectica, 2025. Available at: <https://doi.org/10.48021/978-65-270-6781-8-c13>. Access em: 11 out. 2025.

WILSON, SA **Commentary on: Muscular strength gains during resistance training are attenuated with soy in comparison with latin foods or usual protein intake in older adults: a randomized controlled trial.** *Clinical Nutrition*, Elsevier BV, v. 35, no. 6, p. 1575-1576, 2016. Available at: <https://doi.org/10.1016/j.clnu.2016.07.005>. Access em: 11 out. 2025.

YASUDA, J.; MURATA, K.; HASEGAWA, T.; YAMAMURA, M.; MAEO, S.; TAKEGAKI, J.; TOTTORI, N.; YOKOKAWA, T.; DIE.; ARIMITSU, T.; NISHIKORI, S.; FUJITA, S.
Relationship between protein intake and resistance training–induced muscle hypertrophy in middle-aged women: A pilot study. *Nutrition*, Elsevier BV, v. 97, p. 111607, 2022. Available at: <https://doi.org/10.1016/j.nut.2022.111607>. Access em: 11 out. 2025.

YASUDA, J.; TOMITA, T.; ARIMITSU, T.; FUJITA, S. **Evenly distributed protein intake over 3 meals increases resistance exercise–induced muscle hypertrophy in healthy young men.** *The Journal of Nutrition*, Elsevier BV, v. 150, no. 7, p. 1845-1851, 2020. Available at: <https://doi.org/10.1093/jn/nxaa101>. Access em: 11 out. 2025.