



## THE CLINICAL COMPLEX ARISING FROM THYROID DISORDERS: A GENERAL APPROACH

## THE CLINICAL COMPLEX ARISING FROM THYROID DISORDERS: A GENERAL APPROACH

Tarciany das Neves Pulcino

Medicina by Cesmac

Email:[weto\\_25@hotmail.com.br](mailto:weto_25@hotmail.com.br)

Pedro Gonçalves Bannitz

Medicina, Anhembi Morumbi

[Peedrobannitz@icloud.com](mailto:Peedrobannitz@icloud.com)

Nathan Arenas Périco

Doctor at UniCesumar

Email:[Nathan\\_0010@hotmail.com](mailto:Nathan_0010@hotmail.com)

Ana Karla Alves do Carmo

Medicine, Antônio Carlos Institute of Tocantins (ITPAC Porto)

Email:[Draanakarlaalves7@gmail.com](mailto:Draanakarlaalves7@gmail.com) Solange Mendes Vieira

Pediatrics Resident at the Catholic University of Pelotas UCEPEL

Email:[Solangevieira121@gmail.com](mailto:Solangevieira121@gmail.com) Isabella Frauches Martin

Medicina by UNICESUMAR E-mail:[Isafranches.martin@gmail.com](mailto:Isafranches.martin@gmail.com) Ana

Carolina Alves de Lucena Medicine from UERN

Email:[Ana\\_carolina\\_1990@hotmail.com](mailto:Ana_carolina_1990@hotmail.com)

Arytanna Borges da Silva Castro

General Surgeon at Santa Casa de Misericórdia de São Sebastião do Paraíso/ Faculdade UNIPAC -Juiz de Fora

Email:[arytannaborges@gmail.com](mailto:arytannaborges@gmail.com)

Francisco Alípio de Oliveira Santiago

University Teacher

Email:[alipio.santiago@gmail.com](mailto:alipio.santiago@gmail.com)

Marcus Vinicius Meirelles Rodrigues

Doctor from the Federal University of Roraima (ufrr)

Email:[marcusviniciusmrodrigues@hotmail.com](mailto:marcusviniciusmrodrigues@hotmail.com)

Barbara Yanaisa Castellanos Rodriguez

Higher Institute of Medical Sciences of Santiago de Cuba

Email:[Babiyani89@gmail.com](mailto:Babiyani89@gmail.com) João Pedro Mendonça Raphael

Braz

Medicine by Faculdades de Medicina Nova Esperança/FAMENE Email:

[mraphaelbraz@gmail.com](mailto:mraphaelbraz@gmail.com) Clarissa Verena Ferraz Lisboa Medical,

Uncisal

1

Email:[claraFerraz@hotmail.com](mailto:claraFerraz@hotmail.com)

Gustavo Silva Campos

Medicine by FAMINAS

Email:[lealcampos@gmail.com](mailto:lealcampos@gmail.com)

Rogério Leal Campos

Gastroenterologist at the Faculty of Medical and Health Sciences of Juiz de Fora E-mail:

[Dr.rogeriocampos@gmail.com](mailto:Dr.rogeriocampos@gmail.com) Bruna Gomes Dias Negreiros



Medicine, UEMA

Email: [Brunanegreiros09@gmail.com](mailto:Brunanegreiros09@gmail.com)

Sarah Pereira Lima Xavier

Doctor at the Catholic University of Brasília E-

mail: [sarahplxavier09@gmail.com](mailto:sarahplxavier09@gmail.com) Jessica

Erculano da Silva Medicine – UNCISAL

Email: [jessicaerculano@gmail.com](mailto:jessicaerculano@gmail.com)

Danilo José Brito de Arruda

Doctor – Federal University of Maranhão

Email: [Danillojj12@hotmail.com](mailto:Danillojj12@hotmail.com) Igor Roberto

Guilhon Costa Medicine from UFMA

Email: [igor.quilhon@discente.ufma.br](mailto:igor.quilhon@discente.ufma.br)

Lucas Luiz Fitipaldi Ferreira

Medicine by Centro Universitário de Maringá

Email: [lucasffitipaldi@outlook.com](mailto:lucasffitipaldi@outlook.com) Abraão Pedro

Araújo Almeida Medicine from Universidade

Potiguar Email: [Abraaopedro2015@outlook.com](mailto:Abraaopedro2015@outlook.com)

Wesley Figueiredo de Lucena Medicina, FACISA-

CG PARAÍBA Email: [Figueiredo.wesley@gmail.com](mailto:Figueiredo.wesley@gmail.com)

Giovanna Midory Takeda

Doctor at Nove de Julho University Campus Guarulhos Email

[Midorytakeda@gmail.com](mailto:Midorytakeda@gmail.com) Antonio Emílio Couto Fardin

Doctor from Vila Velha University Email:

[antonioecf99@gmail.com](mailto:antonioecf99@gmail.com) Lucas Rosa

Doctor from Universidade Brasil Email:

[lucasrosa.1994@hotmail.com](mailto:lucasrosa.1994@hotmail.com) Amanda

Ricardi Massochin Doctor at Centro

Universitário INGÁ Email:

[amandaricardi@hotmail.com](mailto:amandaricardi@hotmail.com) Paula

Pessoa Pinheiro

Doctor from the Federal University of Cariri – UFCA

Postgraduate in Teaching in Higher Education in Health, Centro Universitário Cidade Verde – UniCV Email

[Paula.ppsk@gmail.com](mailto:Paula.ppsk@gmail.com) José Rodrigues dos Santos Júnior Doctor by FAHESP/IESVAP Email

[loserodriguesjunior19@gmail.com](mailto:loserodriguesjunior19@gmail.com) Marcos Fernando Neiva de Alencar Lustosa Rocha Medicina by Unifacid

Wyden

Email: [marcosfernandolr1@hotmail.com](mailto:marcosfernandolr1@hotmail.com)

Igor Gabriel Silva Oliveira

Doctor from the Federal University of Jataí

Email: [contato@igorgabriel.com.br](mailto:contato@igorgabriel.com.br)

Amanda Sousa de Andrade Medicine

by PUC campinas Email:

[Amanda.sousa062@hotmail.com](mailto:Amanda.sousa062@hotmail.com) Isabela

Dablan Samara Fazilari Medicine, FMABC

Email: [isafazilari@gmail.com](mailto:isafazilari@gmail.com)

Raphael Dalla Paula Lopes

two



Medicine by Centro Universitário Facisa (UNIFACISA)

Email:[raphadalla13@gmail.com](mailto:raphadalla13@gmail.com) Lucas Marvilla Fraga de Mesquita

Medicine by the Institute of Medical Education – IDOMED

E-mail:[lucasmfdm@gmail.com](mailto:lucasmfdm@gmail.com) Monique Albuquerque Amorim Medicina UNIMA

Email:[Monique\\_albuquerque@hotmail.com](mailto:Monique_albuquerque@hotmail.com)

João Paulo Hammuy

Doctor studying specialization in Family and Community Medicine, UFMG, Belo Horizonte/MG Email:

[joao\\_paulo@outlook.com](mailto:joao_paulo@outlook.com) Isadora Raineri Sgarbi

Medicine by Centro Universitário Barão de Mauá- CUBM

Email:

[Isa.irs@hotmail.com](mailto:Isa.irs@hotmail.com)

Nildo Redivo Junior

Gynecologist and Obstetrician, studying for a doctorate at the Universidade do Oeste Paulista.

email:[nildodr@hotmail.com](mailto:nildodr@hotmail.com) Maria Lorena Ribeiro Redivo

Studying medicine at the Universidade do Oeste Paulista

Email:[marialorenaredivo@gmail.com](mailto:marialorenaredivo@gmail.com) Marcos Vinicius

Dantas Pereira Pamplona Medicina – FCM-PB

Email :[mviniciuspamplona@gmail.com](mailto:mviniciuspamplona@gmail.com)

Rodrigo Alves Giacomelli

Medicine, Unicesumar (graduated in 2023)

[ralvesgiacomelli@gmail.com](mailto:ralvesgiacomelli@gmail.com) Millena

Gonçalves de Carvalho Ferreira Medicina,

UNIDERP- Anhanguera E-mail:

[Millena.qdcb@gmail.com](mailto:Millena.qdcb@gmail.com) Allan Maia de

Jesus

Medicine by the Institute of Medical Education – IDOMED E-

mail:[md.allan@hotmail.com](mailto:md.allan@hotmail.com) Wendryus William De Lima

Medicine, UNISL – São Lucas University – PVHO, RO Luana

Beltrame Zancanela

Doctor from Vila Velha University (ES)

Email:[Luana-zancanela@hotmail.com](mailto:Luana-zancanela@hotmail.com)

Cristiane Bruno de Siqueira Andrade

Medicina UESB

Email:[Crisbruno23@icloud.com](mailto:Crisbruno23@icloud.com) Tarsila

Stockler Ravaschieri Martinolli Medicina –

State University of Pará E-mail:

[tarsilastockler@gmail.com](mailto:tarsilastockler@gmail.com) Samuel Gouveia

da Costa Duarte Doctor by UFPB

Email:[samuelqcduarte@gmail.com](mailto:samuelqcduarte@gmail.com)

3

Nadjila Gabriela Santana Sidani

Medicine, PUC-SP

Email:[nadjilasidani@gmail.com](mailto:nadjilasidani@gmail.com) Vivian de

Carvalho Veleda Moraes Doctor from the

Federal University of Pelotas Email:

[vivenmoraes@live.com](mailto:vivenmoraes@live.com) Tiago Wanderley

Queiroga Lira

Medicine, Faculty of Medicine Nova Esperança da Paraíba (FAMENE)



Email: [twqlira@gmail.com](mailto:twqlira@gmail.com)

Isabelle Laís Oliveira dos Santos Lira  
Medicine, Faculty of Medical Sciences of Paraíba (FCM-PB)  
Email: [Isabelle\\_oliveiras@outlook.com](mailto:Isabelle_oliveiras@outlook.com) Hérica Silva Florentino  
Medicina, UNIMA

Email: [Hericaflorentino11@gmail.com](mailto:Hericaflorentino11@gmail.com)

Vinicius Barbosa Carvalho Medicine,  
Universidade Brasil E-mail:  
[Viniciusbcarvalho090@gmail.com](mailto:Viniciusbcarvalho090@gmail.com) Caroline  
Barros Figueira  
Medicine from the Federal University of Tocantins  
Email: [carolinebarrosfigueira@gmail.com](mailto:carolinebarrosfigueira@gmail.com) Talita  
Maris Campos  
Nutrologist at Monte Pascal University Email:  
[dratalitamaris@gmail.com](mailto:dratalitamaris@gmail.com) Amanda Batista de  
Almeida Medicina, FAMESC

Email: [amandabatistadealm@gmail.com](mailto:amandabatistadealm@gmail.com)

Juliana Marques Amorim  
Medicine by UNIFOR  
Email: [jujumarkes.amorim@gmail.com](mailto:jujumarkes.amorim@gmail.com) Ana  
Carolina Sá de Souza Carvalho Medicine from  
the Federal University of Acre Email:  
[dranacarolinasa@gmail.com](mailto:dranacarolinasa@gmail.com) Gabrielly  
Rodrigues Paniago Medical Resident, Medical  
Clinic Email: [paniagogabrielly@gmail.com](mailto:paniagogabrielly@gmail.com)  
Lucas Camelo Ferreira Faria

Medicine, UniRV – University of Rio Verde – Campus Aparecida de Goiânia  
Email: [Lucascamelofaria@gmail.com](mailto:Lucascamelofaria@gmail.com) Luciana Sousa Simões

Medicine from the Polytechnic and Artistic University UPAP Facultad de Ciencias de La Salud  
Email: [Lucianasousasimoes.ls@gmail.com](mailto:Lucianasousasimoes.ls@gmail.com) Andressa Barros Tenório Nunes de Carvalho  
Medicine from the Faculty of Medicine of Olinda (FMO) Email: [andressatenoriooo@gmail.com](mailto:andressatenoriooo@gmail.com)  
Laura Boldrini Niero

Medicine, Pontifical Catholic University Sorocaba E-  
mail: [Laura.niero@hotmail.com](mailto:Laura.niero@hotmail.com) Ana Carolina Mais  
Araújo Medicine by CESMAC

Email [Ana.carolina.maia.araújo@gmail.com](mailto:Ana.carolina.maia.araújo@gmail.com)

Brian Vicente  
Medicine by the Faculty of Medicine of ABC E-  
mail: [brianfmabc@gmail.com](mailto:brianfmabc@gmail.com)

4

Andressa Campos de Sousa  
Uninovafapi Medicine, Teresina, Piauí  
[dressacsousa@gmail.com](mailto:dressacsousa@gmail.com)

Valeria Coelho Denipote Barberi  
Medicine from Anhembi Morumbi University – Campos SP  
Email: [Leladenipote@gmail.com](mailto:Leladenipote@gmail.com) Lucas Arcoverde Teófilo

Medicine by the University of Fortaleza  
Email: [lucas\\_teoarc@hotmail.com](mailto:lucas_teoarc@hotmail.com)



Mateus Arcoverde Teófilo

Medicine by the Federal University of Cariri

Email: [mateusatf@gmail.com](mailto:mateusatf@gmail.com) Beatris Tres

Medicine by UNINTER

Email: [beatris.tres@hotmail.com](mailto:beatris.tres@hotmail.com)

Ana Carolina Munuera Pereira

Medicina, UNOESTE

Email: [Aninha\\_munuera@hotmail.com](mailto:Aninha_munuera@hotmail.com)

## Summary

The thyroid is an endocrine gland, subject to hormonal stimulation, that is, by the hormone TSH, which is synthesized by the pituitary gland and sent to the thyroid via the bloodstream, thus synthesizing the hormones T4 (thyroxine) and T3 (triiodothyronine), which are responsible for control metabolic functions. Dysfunctions can occur such as hypothyroidism and hyperthyroidism, which are associated with factors such as pregnancy, aging, autoimmune disease and drug interactions with a certain type of drug, all of which can disrupt hormonal levels. Excessive or insufficient production of thyroid hormones can cause pathologies secondary to thyroid dysfunction. Therefore, knowledge about the proper functioning of the thyroid gland is essential, seeking to preserve homeostasis and, subsequently, a better quality of life for the patient. The diagnosis, that in both conditions, serum levels of Thyroid Stimulating Hormone (TSH) and thyroid hormone are essential for this, and treatment are essential for the patient to lead a normal life. The objective was to carry out a literature review on general aspects of the clinical complex arising from thyroid disorders, addressing the respective diagnosis and treatment. **Keywords: Endocrine gland; hormonal disorder; thyroid.**

## Abstract

The thyroid is in the endocrine gland, subject to hormonal stimulation, that is, by the hormone TSH, which is synthesized by the pituitary gland and sent to the thyroid via the bloodstream, thus synthesizing the hormones T4 (thyroxine) and T3 (triiodothyronine), which are responsible for controlling metabolic functions. Dysfunctions can occur such as hypothyroidism and hyperthyroidism, which are associated with factors such as pregnancy, aging, autoimmune disease and drug interactions with a certain type of drug, all of which can disrupt hormonal levels. Excessive or insufficient production of thyroid hormones can cause pathologies secondary to thyroid dysfunction. Therefore, knowledge about the proper functioning of the thyroid gland is essential, seeking to preserve homeostasis and, subsequently, a better quality of life for the patient. The diagnosis, that in both conditions, serum levels of Thyroid Stimulating Hormone (TSH) and thyroid hormone are essential for this, and treatment are essential for the patient to lead a normal life. The objective was to carry out a literature review on general aspects of the clinical complex arising from thyroid disorders, addressing the respective diagnosis and treatment.

**Keywords: Endocrine gland; hormonal disorder; thyroid.**

## Introduction

5 The thyroid gland, similarly known as the butterfly gland due to its shape, is one of the largest glands in the human endocrine system. Located in the anterior portion of the neck. The thyroid is essential for our lives, as it is responsible for regulating our body's metabolism. Acting on various organs and functions of the body, throughout life. It influences organs such as the heart, brain, liver and kidneys; helps regulate the growth and development of the body; menstrual cycles and female fertility; It is important for memory, concentration, mood and emotional control. All of these events occur through the thyroid hormones produced by the gland: triiodothyronine, known as T3, and thyroxine, known as T4. These hormones, when entering the bloodstream, reach any tissue or organ and balance various functions. These hormones cannot become unbalanced, as their increase or decrease causes negative reactions in the body, such as hypothyroidism and hyperthyroidism.

The following article aimed to describe in a narrative way considerations regarding the thyroid, as well as



as disorders that cause pathologies, making early diagnosis and therapy essential.

## Methodology

This is a qualitative narrative review study, appropriate to describe the clinical complex arising from thyroid disorders. It consists of a broad analysis of the literature, without establishing a rigorous and replicable methodology at the level of data reproduction and quantitative answers to specific questions).

As it is a bibliographic analysis on the theory of mind and the understanding of this competence in the adult individual, articles indexed in the SciELO, PubMed, Latindex, Google Scholar databases were retrieved, during the month of January 2024, using the reference period as the last 5 years.

The index terms or descriptors thyroid, adrenal gland, hormonal pathologies were used, isolated or in combination, without delimiting a temporal interval. The criteria used for inclusion of publications was to have the expressions used in the searches in the title or keywords, or to have it explicit in the abstract that the text is related to the association of thyroid disorders and linked pathologies. The excluded articles did not meet the established inclusion criteria and/or were duplicated, that is, publications retrieved from more than one of the databases. Dissertations and theses were also excluded.

After the target information was retrieved, the titles and abstracts were initially read, with no publications being excluded at this stage. Subsequently, the complete reading of the 31 texts was carried out. As axes of analysis, we initially sought to classify the studies according to the particularities of the sampling. From there, the analysis of the theoretical foundation of the studies continued, as well as the observation of the general characteristics of the articles, such as year of publication and language, followed by their objectives. Finally, the methodology applied, results obtained and discussion were assessed. Specifically, to analyze the scientific production identified, specific qualitative and/or quantitative data processing techniques were not used, and each text was analyzed.

## Results and discussion

The search for articles that made up this study identified 77 references on thyroid disorders in the aforementioned databases, of which 40 publications were included in the review. Among the selected studies, 38 articles present a theoretical approach and 02 are case studies. The prevalence of publications in English was observed, representing 84% of the total, when compared to Spanish (9.6%) and Portuguese (6.4%).

From a careful reading of the selected texts, it was possible to observe that the studies are unanimous regarding the initial contextualization of the thyroid being responsible for the production of the hormones tetraiodothyronine or thyroxine (T4) and tri-iodothyronine (T3). Both originate from the complement of iodine radicals to tyrosine residues contained in a large 660 kDa glycoprotein, called thyroglobulin (Tg), secreted by follicular cells and stored within the thyroid follicle. Tg corresponds to 70 to 80% of the protein content of the thyroid.

The uptake of iodine from the bloodstream is actively carried out by the Na/I Symporter (NIS) protein, present in the basement membrane of the follicular cell. The enzyme responsible for the oxidation of iodine ions and their binding to Tg is thyroperoxidase (TPO), present in the apical membrane of follicular cells. Initially, molecules composed of a tyrosine residue, linked to Tg, and an iodine molecule — monoiodotyrosine (MIT) — or two iodine molecules — diiodotyrosine (DIT) are formed. MIT and DIT are cleaved and released from Tg by the action of TPO, phagocytosed and linked, so that two DITs form T4 and one DIT, added to an MIT, forms T3

6

T3 and T4 circulate in plasma bound to carrier proteins, such as thyroxine-binding globulin (TBG), and albumin. Only a tiny fraction of these hormones — 0.04% of T4 and 0.4% of T3 — circulates in plasma in free form, not bound to proteins, and is responsible for biological activity of thyroid hormones.

The regulation of thyroid function is mediated mainly by the pituitary hormone thyrotropin, or TSH, which stimulates the synthesis and release of hormones, as well as thyroid growth. TSH, in turn, is positively regulated by hypothalamic thyrotropin-releasing hormone (TRH) and negatively regulated by somatostatin. Secreted T3 binds to receptors in the hypothalamus and pituitary gland, where it inhibits secretion

of TRH and TSH by negative feedback, thus regulating thyroid function

T4 is predominant in thyroid secretions, but T3 is responsible for most of the gland's biological action. Justified by the fact that T3 has an affinity for the thyroid hormone receptor four to 10 times more potent than T4

Intracellular T3 has two main sources: synthesis from T4 and secretion by the thyroid itself.

Deiodination: extraction of an iodine atom from T4, making it T3, occurs through the action of enzymes called 5'-deiodinases, present in practically all organs and systems.

In fact, the peripheral conversion of T4 to T3 is responsible for the majority of circulating T3. T3 binds to nuclear receptors belonging to the superfamily of retinoid receptors, present in practically all organs, regulating the transcription of specific gene sequences, which will lead to the most diverse biological actions of the hormone.

The prevalence of hypothyroidism varies in different regions of the world, being more common in places where there is a deficiency or excess of iodine in the diet, despite being common in iodine-replete areas. It is about 10 times more common in females. Congenital hypothyroidism affects 1:4,000 newborns, and central hypothyroidism is found in 1:20,000 individuals.

Hypothyroidism is one of the most common endocrine pathologies, mostly triggered by deficient synthesis of thyroid hormones.

The etiologies of hypothyroidism can be grouped into:

- Primary: conflict located in the peripheral gland — Central thyroid disease
- Secondary: problem located in the pituitary gland — TSH deficiency;
- Tertiary: problem located in the hypothalamus — TRH deficiency.

| Primary                      | Secondary         | Tertiary  |
|------------------------------|-------------------|---|
| Iodine deficiency            | Pituitary adenoma | Hypothalamic dysfunction                          |
| Hashimoto's thyroiditis      |                   | Idiopathic  |
| Excess iodine                |                   | Surgery or irradiation of the hypothalamic region |
| Defects in hormone synthesis |                   |   |

The abundance of hypothyroidism episodes, representing 95%, is primary. In Brazil, the main cause of this syndrome is chronic autoimmune thyroiditis, or Hashimoto's thyroiditis. Subacute granulomatous, lymphocytic and postpartum thyroiditis can also lead to hypothyroidism, which, in these situations, is usually transient

Cretinism is a mental disability caused by **congenital hypothyroidism**, during the development of the newborn. During this period, the absence of the hormone thyroxine disrupts brain maturation.

The main cause of the disorder is a defect in the **thyroid formation**, but it is also possible that it occurs due to an enzyme deficiency during the development of the hormone. There is one case of cretinism every three thousand births and it is possible to identify the disease through **Foot test**.

Myxedema is usually caused by prolonged hypothyroidism, myxedema is a hard edema with an opaque skin appearance. The disorder occurs five times more frequently in women than in men. People with myxedema experience edema on the face and eyelids, forming "bags" under the eyes. There is also an accumulation of proteins produced in hypothyroidism.

## Hashimoto's thyroiditis

7

Chronic autoimmune thyroiditis — most common autoimmune disease

Goiter is common in around 90% of cases (goiterogenic form). The volume expansion of the Rheoid is generally mild to moderate, symmetrical, slow to develop over decades, and firm in consistency, like rubber, on palpation. Ten percent present with a diminished thyroid, perhaps reflecting the final stage of the gland's chronic destructive process — the atrophic form.

Hashimoto's thyroiditis can, therefore, appear in an incipient form, without hypothyroidism itself and with a normal-sized, enlarged or diminished thyroid. If you have Hashimoto's thyroiditis,

Progression from euthyroidism to subclinical hypothyroidism and then to frank hypothyroidism occurs at a rate of 3 to 15% per year.

The vast majority have high serum titers of antithyroid antibodies, mainly anti-TPO, present in 80 to 99% of cases, and anti-Tg, present in 30 to 60% of cases. However, they are not specific, as they can be elevated in people without thyroid disease — anti-Tg, in up to 10% of the general population, and anti-TPO, in up to 15%.

Hypothyroidism occurs due to immune-mediated destruction of the thyroid parenchyma. Thyroid autoimmunity is of the cytotoxic type, full of cytotoxic or natural killer “T” cells

Diagnosis is based on the abundant identification of autoantibodies (anti-TPO or anti-Tg) in a person with goiter and/or hypothyroidism. Furthermore, the hypoechogenicity of the thyroid parenchyma on ultrasound. The anatomopathological examination may reveal lymphocytic infiltration.

## Thyroid nodules

They correspond to benign or malignant tumors in the thyroid gland, mostly proportional to the increase in age. The predominance is benign. The etiology includes hyperplastic colloid goiter, thyroid cyst, thyroiditis, thyroid adenoma

The nodules may be painful or asymptomatic. Pain indicates thyroiditis or hemorrhage into a cyst. The asymptomatic nodule may be malignant, but is usually benign

The clinical picture of hyperthyroidism, usually palpitations, heat intolerance, weight loss, tremors leads to the hypothesis of hyperfunctional adenoma or thyroiditis. Hypothyroidism, which leads to intolerance to cold, causes weight gain, fatigue, suggests Hashimoto's thyroiditis

Predisposing factors for thyroid cancer include a history of thyroid radiation, especially in childhood, male gender, previous family history of thyroid cancer or multiple endocrine neoplasia type 2, solitary nodule, dysphagia, dysphonia. Increase in size (particularly rapid growth or growth while undergoing thyroid hormone suppression treatment)

On physical examination the symptoms that suggest thyroid cancer include stony stiffness or fixation to adjacent structures, cervical lymphadenopathy, and hoarseness (due to recurrent laryngeal nerve palsy).

The initial evaluation of a thyroid nodule is performed with tests for Thyroid Stimulating Hormone (TSH) and anti-thyroid peroxidase antibodies. If thyroid-stimulating hormone (TSH) is low (consistent with hyperthyroidism), a radioactive iodine test is performed. Nodules with increased iodine uptake (warm) are rarely malignant.

If thyroid function tests do not indicate hyperthyroidism or Hashimoto's thyroiditis, the next step is to examine the nodule with a thyroid ultrasound. Ultrasonography is useful for determining the size of the nodule, but may not be able to distinguish between benign and malignant lesions. Thyroid cancer is suggested in case of hypoechogenicity of the nodules; accentuated internal vascularization; uneven edges; nodule height greater than its width; irregular macrocalcifications; fine dotted psammomatous microcalcifications (papillary-type thyroid carcinoma); or dense homogeneous irregular calcification (medullary thyroid carcinoma).

For nodules with suspicious characteristics, fine needle aspiration biopsy (FNAB) performed under ultrasound guidance is the procedure of choice and the most economical means of distinguishing benign from malignant nodules. In general, FNAC is not recommended for nodules <1 cm or nodules that are entirely cystic.

The cytological examination of cells obtained from FNAC is capable of distinguishing benign from malignant lesions. For cytologically indeterminate lesions, molecular analysis of FNA aspirate looking for mutations and rearrangements or measuring calcitonin looking for possible medullary carcinoma can add clarity sufficient diagnostics to enable an optimally informed treatment recommendation.

Treatment of a thyroid nodule is based on therapy for the underlying disease. Nodules in patients with Hashimoto's thyroiditis or other causes of goiter associated with hypothyroidism may stabilize or regress with thyroxine replacement at doses that make the patient euthyroid. For small benign nodules not associated with hypothyroidism, TSH suppression with thyroxine effectively shrinks the nodule in no more than half of cases, and is rarely done. Thyroxine is not used to treat neoplastic nodules.

The size of large benign nodules that compress adjacent cervical structures may decrease.

nir after radiofrequency ablation. Partial or complete thyroidectomy is effective in treating nodules that continue to grow or cause compressive symptoms.

Toxic nodules can be treated with radioactive iodine to control hyperthyroidism and reduce their size. Nodules that are not due to thyroid cancer, do not cause compressive symptoms, and are not associated with hypothyroidism or hyperthyroidism can be safely monitored with periodic physical examinations of the neck and, if growth of the nodules is suspected, by repeat ultrasound evaluation.

### Goiter

Goiter, also known as puffiness, is an increase in volume in the thyroid. Normally, it is associated with hypothyroidism or hyperthyroidism, but this association is not necessary or mandatory.

Many goiters have nodules and are called nodular goiters, which do not necessarily interfere with thyroid function and therefore do not generate symptoms of hypo or hyperthyroidism.

Diffuse goiter: May occur in thyroid dysfunction or when there is an iodine deficiency.

Nodular goiter: It can be uninodular (just one nodule) or multinodular (more than one goiter).

Nodular goiter is most often diagnosed after age 50. The patient may have had the goiter for several years and only detect it when compressive symptoms such as dysphagia and hoarseness appear; or even just diagnose because it has become hyperfunctioning, causing hyperthyroidism. Another episode that may occur is presenting the diagnosis only when investigating associated hypothyroidism.

In addition to palpation, blood tests and ultrasound must be ordered by the doctor. To assess whether there is a change in thyroid function, TSH measurement is the best initial investigation test.

Further investigation and treatment will depend on the cause, symptoms, changes (or not) in the functioning of the gland and the characteristics of the nodules on ultrasound (in cases of nodular goiters).

### Conclusion

It was found that the thyroid plays a very important role in the patient's life. The hormones secreted by it and stimulated by TSH, as long as they are at normal levels, are responsible for controlling various functions of the body. It is worth highlighting the importance of carrying out routine tests to check whether the levels of these hormones are properly regulated, as their dysfunctions can cause harm to the patient's daily life, especially in women over 40 years of age, the elderly and pregnant women.

### References

1. BARROS, ACS; et al. Biochemical pharmacist: an approach focused on TSH and diseases of the thyroid. Health and Human Development, [s.l], v. 6, no. 1, p.67-67, 1 mar. 2018. La Salle University Center – UNILASALLE.
2. SOUZA, LSL; et al. Analysis of iodine nutritional status in hypertensive and non-hypertensive pregnant women in Salvador-Bahia. Journal of Medical and Biological Sciences, Salvador, v.14, n.3, p.323-330, 2015.
3. FONSECA, CW; MELEK, FE Drugs widely used in clinical practice that interact with thyroid hormones. Brazilian Society of Clinical Medicine Magazine, p. 307-313, February 5, 2014.
4. LIMA, J. et al Thyroid dysfunction and amiodarone. Brazilian archives of endocrinology and metabolism, February 2013.
5. MALHOTRA, S.; et al. Topical Iodine-Induced Thyrotoxicosis in a Newborn with a Giant Omphalocele. American Journal Of Perinatology Reports, [s.l], v. 06, no. 02, p.243-245, 28 jun. 2016. Georg Thieme Verlag KG.
6. OLIVEIRA, V.; MALDONADO, RF Hypothyroidism and hyperthyroidism – a brief review of thyroid dysfunctions. Revista Interciência & Sociedade, vol.3, N.2, page. 36-44, 2014.
7. OLIVEIRA, RA Differentiated thyroid cancer, diagnosis and treatment with iodine-131. Radiology Technology Course, Interdisciplinary Research Center – 2015.



8. PEDROSO, B.; TANEDA, M. Analysis of cases of hypothyroidism in the northwest region of Mato Grosso do Sul. *AJES Health Magazine*, 2014.
9. SANTOS, L.M.; et al. Temporal evolution of mortality from thyroid cancer in Brazil from 2000 to 2012. *Revista Brasileira de Análises Clínicas*, p.133-137. Approved March 10, 2016.
10. SILVA, AIN Maternal subclinical hypothyroidism during pregnancy and consequences in school-aged children. *Bibliographic Review Article – Integrated Masters in Medicine*, 2017.
11. SÚSS, SK A de. Clinical outcomes in patients with differentiated thyroid cancer treated with total thyroidectomy and 30 mCi of radioactive iodine versus patients treated with total thyroidectomy alone Curitiba, 2017.
12. TAVARES, MB; et al. A. Amiodarone-induced thyroid function disorders. *Salus Journal Of Health Sciences*, pg. 39-47, March 15, 2016.
13. TOMAZ, FD D; et al. Prevalence of Hypothyroidism in the Elderly in the Municipality of Taubaté-SP. *Brazilian Journal of Health Sciences* vol.20, n.3, pages 235-240, 2016.
14. Pinheiro, VP & Nunes, CP. (2019). Therapeutic management in hypothyroidism and pregnancy. *Journal of Family Medicine and Mental Health*.
15. Ríos, A, Rodríguez, JM, Febrero, B, Acosta, JM, Torregrosa, N, Balsalobre, M & Parrilla, P. (2011). Histological and immunohistochemical profile of sporadic and familial medullary thyroid carcinoma. *Endocrinology and Nutrition (English Edition)*, 58(10), 521-528.
16. Rodrigues, BM, Toledo, KA & Nogueira, CR. (2015). Human body: several interconnected systems. *Learning Science*, 4(1): 79-83.
17. Rosário, PW, Ward, LS, Carvalho, GA, Graf, H, Maciel, R, Maciel, LMZ & Vaisman, M. (2013). Thyroid nodule and differentiated thyroid cancer: update of the Brazilian consensus. *Brazilian Archives of Endocrinology & Metabology*, 57(4), 240-264.
18. Silva, RV, Alves, LCB, Doro, RB & Malthez, ALMC. (2019). Assessment of Absorbed Dose in the Thyroid in Mammography Exams. *Brazilian Journal of Medical Physics*, 13(3), 24-27.
19. Tavares, MB, Motta, PRV, Barros, VF, Cezana, C, Ferreira, LB, Saar, SMA & Brandão, CDG. (2016). Amiodarone-induced thyroid function disorders. *Revista Salus, Journal of health sciences*, 2(2), 39-47.
20. Tomaz, FDD, Silva, AM, Bissoli, CF, Ferreira, RCA & Fernandes, WS. (2016). Prevalence of Hypothyroidism in the Elderly in the Municipality of Taubaté-SP. *See. Bras. Health Science*, 235-40.