Machine Translated by Google
RCMOS – Multidisciplinary Scientific Journal The Knowledge.
ISSN: 2675-9128. São Paulo-SP.

**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

# Applied Neuroanesthesia and Surgical Rehabilitation in Leprosy: Clinical Advances in Peripheral Neurolysis and its Intersection with Occupational Health in Anesthetic Practice

*Applied Neuroanesthesia and Surgical Rehabilitation in Leprosy: Clinical Advances in Peripheral Neurolysis and the Intersection with Occupational Health in Anesthetic Practice*

Applied neuroanesthesia and surgical rehabilitation in leprosy: clinical advances in peripheral neuroplasia and the intersection with occupational health in anesthetic practice

**Dr. Janaína Leila Batista de Oliveira ,**

Anesthesiologist, Specialist in Neurosciences (Unyleya), Specialist in Occupational Medicine (UnB), Specialist in Medical Residency Preceptorship (Oswaldo Cruz), Medical Residency in Anesthesiology (FEPECS / ESCS), Graduated in Medicine (UFRN).

**Summary**

Leprosy remains a significant public health challenge in many regions of the world, with peripheral nervous system impairment being one of the main causes of functional disability associated with the disease. In this context, decompressive neurolysis procedures play a relevant role in preserving neurological function and rehabilitating patients affected by peripheral neuropathies. This article reviews the scientific evidence related to neuroanesthesia applied in these procedures, exploring its interface with neurophysiological monitoring, occupational medicine, and the neurosciences of human performance.

Beyond the technical aspects involved in anesthetic management, this article discusses how cognitive, emotional, and occupational factors influence decision-making in highly complex surgical environments. The literature reviewed demonstrates that elements such as situational awareness, crisis management, stress regulation, structured communication, and prevention of professional fatigue have a direct impact on patient safety and the performance of perioperative teams.

The analysis shows that the integration of neuroanesthesia, occupational health, and cognitive neuroscience broadens the understanding of the factors that underpin highly reliable surgical systems. The knowledge produced in these areas provides a basis for the development of structured training methodologies, risk management protocols, and professional development models that can be replicated in different healthcare settings. In this way, contemporary anesthetic practice is understood not only as technical support for the surgical act, but also as a strategic field for building organizational cultures focused on safety, operational resilience, and excellence in care.

**Keywords:** Neuroanesthesia; Leprosy; Peripheral Neurolysis; Occupational Health; Applied Neurosciences; Perioperative Safety.

**Abstract**

Leprosy constitutes a chronic endemic disease that promotes peripheral nervous system impairment, requiring decompressive neurolysis surgeries to mitigate physical deformities and neuropathic pain. This review article analyzes the contemporary scientific evidence underlying the practice of applied neuroanesthesia in peripheral neurolysis procedures. The impact of specialized medical volunteering in rehabilitation workshops is explored, and the occupational health determinants for anesthesiologists are discussed in the context of occupational medicine.

Additionally, the application of neurosciences in optimizing clinical performance and decision-making under stress in the surgical environment is analyzed. The results indicate that the association of ultrasound-guided regional anesthesia with neurophysiological monitoring preserves intraneural perfusion and improves functional prognosis. It is concluded that

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google
RCMOS – Multidisciplinary Scientific Journal The Knowledge.
ISSN: 2675-9128. São Paulo-SP.

**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

Investing in the doctor's occupational health and crisis management methodologies in the operating room ensures the sustainability and biological safety of high-complexity anesthetic acts.

**Keywords:** Neuroanesthesia; Leprosy; Peripheral Neurolysis; Occupational Health; Crisis Management.

## Introduction

Leprosy remains a chronic endemic disease with a significant impact on public health.
in developing countries, characterized by the tropism of *Mycobacterium leprae* for
peripheral nervous system. Bacterial invasion of Schwann cells triggers a
A complex immune-mediated inflammatory process that culminates in demyelination and fibrosis.
Perineural and mechanical entrapment of nerve trunks. This progressive neurological damage
It is the main cause of physical disabilities, anatomical deformities, and pain.
Chronic neuropathic conditions that stigmatize affected individuals.
Surgical decompression treatment, implemented through the peripheral neurolysis technique,
It plays a crucial and complementary role to conventional polychemotherapy in prevention.
and in the rehabilitation of neurological sequelae. Surgical decompression aims to restore blood flow.
Intraneural blood flow, relieve tissue pressure, and interrupt the ischemic cascade that destroys tissue.
healthy axons. The ultimate success of this reconstructive intervention depends directly on the healthy axons.
conditional on the surgeon's technical precision and specialized neuroanesthetic management during
the intraoperative phase.
In the context of neurolysis procedures, applied neuroanesthesia takes on several responsibilities.
Scientific interventions that go far beyond the mere provision of immobility and analgesia. The anesthesiologist
pharmacological techniques should be selected that aim to preserve nerve perfusion.
In cases of compromised peripherals, avoid severe hemodynamic fluctuations and allow for the performance of...
Intraoperative neurophysiological monitoring tests. The combination of general anesthesia
Balanced peripheral nerve blocks guided by ultrasound are emerging as
a highly effective strategy.
Performing these surgical procedures in the context of humanitarian actions and workshops.
High-turnover surgical procedures impose severe logistical and operational challenges on the team.
health is involved. The medical professional must demonstrate high technical adaptability and
Psychological resilience to ensure the biological safety of patients in high-risk environments.
Healthcare pressure. This scenario of vocational dedication and medical volunteerism highlights
the social impact of specialized medicine in restoring functionality and dignity of

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google
RCMOS - Multidisciplinary Scientific Journal The Knowledge.
ISSN: 2675-9128. São Paulo-SP.

**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

Neglected populations.

This review article aims to analyze the contemporary scientific evidence regarding from the intersection between applied neuroanesthesia and peripheral neurolysis techniques in leprosy. and the determinants of occupational health for anesthesiologists. The following are discussed: pathophysiological mechanisms of nerve damage, anesthetic strategies that optimize the neurological outcomes, the role of neuroscience applied to clinical decision-making under Stress and the ergonomic and psychosocial risks associated with intensive perioperative practice. in the surgical center.

## Chapter 1: Pathophysiology of Nerve Damage in Leprosy and the Surgical Approach Decompression

Mycobacterium *leprae* has a unique biological affinity for Schwann cells. that form the myelin sheath of the nerves in the human peripheral nervous system. The connection Bacterial transmission occurs through specific endothelial surface receptors, triggering a cellular or humoral immune response that varies according to the immunogenetic profile of the host. This chronic inflammatory process results in severe edema and increased progressive increase in hydrostatic pressure in rigid nerve compartments. Increased intraneural pressure compromises the capillary microcirculation of the vasa nervorum. leading to persistent tissue ischemia and chronic axonal hypoxia in the affected limb. If the If mechanical compression and ischemia are not reversed promptly, the proliferation of Fibrous connective tissue replaces functional nerve fibers with perineural scars. dense. This architectural destruction manifests clinically through sensory deficits and Motor impairment, loss of vital reflexes, and development of claws on the extremities. The surgical decompressive approach, represented by external and internal neurolysis, It is based on the opening of the epineurium and the release of fibrous adhesions that trap the nerve in narrow anatomical channels. The main nerve trunks affected are The nerves that require intervention are the ulnar, median, common fibular, and posterior tibial nerves. Immediate decompression restores the capillary perfusion gradient, allowing for regeneration of the... viable axons and halting the progression of disabling physical deformities. Determining the ideal surgical time is a critical factor in the prognosis of Functional recovery of leprosy patients within the healthcare system. Medical literature. Surgical intervention is recommended in cases of neuritis that do not respond to treatment.

Clinical intervention with corticosteroid therapy, after four to six weeks, in the presence of intractable neuropathic pain.

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google
RCMOS – Multidisciplinary Scientific Journal The Knowledge.
ISSN: 2675-9128. São Paulo-SP.

**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

or when there is a progressive and rapid worsening of sensory or motor functions. The procedure
Early onset of neurolysis is directly correlated with higher reversal rates.
The success of peripheral neurolysis transcends the isolated surgical act, integrating itself into a broader context.
a comprehensive program for prevention and post-operative physical rehabilitation, of a general nature
multidisciplinary. The surgeon and the anesthesiologist work in perfect harmony to
to ensure that the procedure takes place under conditions of tissue stability. The restoration of
Physiological nerve conduction and relief of chronic neuropathic pain are the primary outcomes.
which validate the effectiveness of the decompressive surgical approach as a tool for
social reintegration.

**Chapter 2: Advanced Neuroanesthesia Techniques in Peripheral Nerve Neurolysis**

Selecting the ideal anesthetic strategy for performing peripheral neurolysis in patients
Leprosy requires a careful analysis of the pathophysiological conditions of the nerves.
The factors affected and the extent of the procedure. Modern neuroanesthesia advocates the use of
approaches that combine systemic hemodynamic stability with excellent blockade.
Autonomic and local sensory. Regional anesthesia, specifically plexus blocks or
The use of isolated peripheral nerves has become established as a highly effective and safe modality.
for these interventions.
The advent of ultrasound revolutionized the practice of contemporary regional anesthesia.
allowing direct, real-time visualization of nerve structures, from the blocking needle
and the dispersion of the local anesthetic into adjacent tissues. In nerves affected by
leprosy, which often presents with severe thickening and anatomical distortion.
Due to fibrosis, pre-procedural ultrasound mapping is essential.
This technology minimizes the risk of accidental intraneural injection, a complication that could
exacerbate pre-existing axonal damage.
The choice of local anesthetic drugs is based on the estimated duration of the neurolysis surgery.
and in the need for prolonged postoperative analgesia. Long-acting agents and with
Drugs with lower cardiotoxicity potential, such as ropivacaine and levobupivacaine, are widely...
preferred in concentrations that allow effective sensory blockade without inducing a blockade.
Excessively prolonged motor function. Fine control of the dose and volume of local anesthetic.
It prevents temporary mechanical compression of the peripheral nerve.
When the surgical decompression procedure involves multiple limbs or requires a

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google
RCMOS – Multidisciplinary Scientific Journal The Knowledge.
ISSN: 2675-9128. São Paulo-SP.

**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

long-term approach, the combination of regional anesthesia with intravenous general anesthesia.
Total emerges as a synergistic technique of excellence. The continuous infusion of propofol and
remifentanil, guided by brain depth monitoring, provides excellent
control of the autonomic response to surgical stress. This modality preserves the
self-regulation of blood flow in peripheral tissues and ensures a gentle awakening of
patient.

Intraoperative management requires constant vigilance from the anesthesiologist regarding...
patient positioning on the operating table, in order to avoid stretching or strain injuries
Secondary compression of healthy or weakened nerves. The use of protective cushions and the
Strict anatomical alignment of the limbs are mandatory preventive measures during
The entire procedure. Administering high-precision neuroanesthesia ensures optimal operating conditions.
ideal for reconstructive microsurgery, which is directly reflected in the acceleration of
neurological rehabilitation.

**Chapter 3: Medical Volunteering and Humanitarian Actions in the Rehabilitation of Endemic Diseases Chronicles**

The undertaking of medical volunteer work and humanitarian missions represents a
an essential dimension of the social responsibility of specialized medicine in the face of disease.
Neglected problems that plague economically vulnerable populations. Prevention workshops
and physical and surgical rehabilitation, structured in endemic regions to provide care to
Patients with leprosy function as centers for concentrating medical technology and...
High-impact clinical expertise. These initiatives aim to reduce long waiting lists for
procedures in the public healthcare system.

The participation of the anesthesiologist as a volunteer in these high-risk campaigns.
Turnover demands skills that go beyond the usual technical excellence.
in large, stable hospital centers. The professional must demonstrate high adaptability.
operational for managing the intense flow of procedures in outpatient settings or
in surgical centers with varying logistical resources. Optimizing operating room times, the
Standardization of regional anesthesia protocols and rapid pre-anesthetic screening are
determinants for safety.

The humanitarian impact of performing peripheral neurolysis in a surgical campaign setting.
It is consolidated in the restoration of functional and work capacity to individuals who...
They were marginalized due to the physical deformities resulting from leprosy.

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google
RCMOS - Multidisciplinary Scientific Journal The Knowledge.
ISSN: 2675-9128. São Paulo-SP.

**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

to surgically relieve chronic neuropathic pain that prevents sleep and work, the
Excellent medical intervention restores the patient's dignity and autonomy. The experience
The clinical skills acquired in these workshops enrich the professional's humanistic and technical knowledge.
Interprofessional coordination and cooperation between different medical specialties, such as
reconstructive plastic surgery, orthopedics, anesthesiology, as well as the teams of
Nursing and physiotherapy are the key to the operational success of the actions.
Voluntary. Prior alignment of goals and synchronized execution of tasks reduce the
The incidence of organizational failures in high-pressure surgical care environments. This
Synergy demonstrates that science-based volunteering has the capacity to replicate patterns.
high.

Specialized medical volunteering also acts as a catalyst for strengthening...
Local health policies and the training of professionals residing in the regions served. A
Transfer of practical knowledge in neuroanesthesia and surgical techniques.
Decompression therapy leaves a lasting technical legacy in the communities it serves. In this way,
Humanitarian actions transcend one-off assistance, taking on the form of...
Health education and well-being promotion strategies.

**Chapter 4: Occupational Medicine and Occupational Health in the Perioperative Context**

The operating room environment is recognized by specialized scientific literature as a
of the most complex and demanding work ecosystems for occupational health of
medical professionals. The anesthesiologist is chronically exposed to
a constellation of physical, chemical, biological, and psychosocial risks in the exercise of their duties.
daily activities. A detailed analysis of these factors from the perspective of occupational medicine is
indispensable for developing effective preventive strategies in the hospital environment.
Occupational exposure to residual anesthetic gases resulting from leaks in systems
The ventilation or exhalation of patients in the recovery room represents a chemical risk.
Silent and persistent. Although modern operating rooms have exhaust systems.
Active, chronic inhalation of traces of nitrous oxide and halogenated agents has been associated
to potential long-term cellular biological effects. Regular quality monitoring
Air quality control and preventive maintenance of equipment are occupational engineering measures.
mandatory.
Physical risks clearly include exposure to ionizing radiation during

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google
RCMOS - Multidisciplinary Scientific Journal The Knowledge.
ISSN: 2675-9128. São Paulo-SP.

**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

Fluoroscopy-guided surgical procedures, common in orthopedics and neurosurgery,
In addition to the inadequate ergonomic workload in the operating room, the anesthesiologist remains...
long periods in static postures or prolonged standing to monitor
vital parameters and perform high-precision procedures. This biomechanical overload
predisposes to the development of work-related musculoskeletal disorders.
In the realm of psychosocial risks, the intrinsic nature of anesthesiology, marked by
need for uninterrupted surveillance and direct responsibility for the patient's life in
Emergency scenarios generate a high mental load and acute stress. Working under emergency conditions...
Frequent night shifts alter the physician's circadian rhythm, resulting in sleep deprivation and
chronic sleep fatigue. This scenario of continuous pressure acts as a powerful trigger for
The development of Burnout Syndrome in the medical field.
The development of robust occupational health programs in hospital institutions and
Specialized clinics are a legal and ethical requirement to mitigate burnout.
Early professional development. Preventive strategies based on cognitive ergonomics, in
rational rotation of shift schedules, ensuring adequate rest breaks and in
Institutionalized psychological support reduces absenteeism from work. Preserving the health of
Having an anesthesiologist is a fundamental requirement to ensure care.

**Chapter 5: Neuroscience Applied to Clinical Performance and Decision Making**

Clinical decision-making in crisis situations in the operating room constitutes one of the
more demanding cognitive processes in medical practice, requiring instant integration.
of sensory data, prior knowledge, and emotional control. The neurosciences applied to
Human performance studies investigate the brain mechanisms that govern human behavior.
Anesthesiologist under conditions of acute stress and life-threatening patient.
Understanding how neural systems work allows for optimizing mental acuity in
perioperative environment.
Under conditions of high stress, massive activation of the amygdala triggers a
autonomic fight-or-flight response, mediated by the sympathetic nervous system and the release
adrenaline and cortisol. This state of heightened alertness, while beneficial for physical reactions.
primitive, it can trigger the phenomenon of emotional hijacking, compromising the functions
executive functions of the prefrontal cortex. In clinical practice, this can manifest itself through
narrowing of the attention span and slowing of reaction time in response to

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google
RCMOS - Multidisciplinary Scientific Journal The Knowledge.
ISSN: 2675-9128. São Paulo-SP.

**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

serious incidents.

Maintaining situational awareness is the primary cognitive defense against error.
Judgment in the management of perioperative crises. Situational awareness is divided into
three fundamental neurocognitive levels: the perception of environmental elements, the
understanding the meaning of this real-time data and projecting the future state of
The patient will be there for the next few minutes. The high-performing anesthesiologist uses heuristics.
structured to avoid overloading working memory during the operating process.
Resource management methodologies in medical crises, inspired by aviation.
commercially, they are based on findings from neuroscience to standardize behaviors that
These guidelines mitigate human error and advocate the use of cognitive checklists.
Speaking aloud, closed-loop communication, and clear task distribution within the team.
multidisciplinary. By externalizing decision-making through visual protocols, it reduces-
if the consumption of metabolic energy in the brain is reduced, while preserving analytical capacity.
The incorporation of emotional regulation and neurocognitive resilience training into
The training of anesthesiologists consolidates the pursuit of high performance and safety.
assistance. The use of prior visualization techniques, breathing control exercises for
Modulation of vagal tone and repeated debriefing sessions after critical events help to
Brain plasticity creates more efficient synaptic pathways. The fusion between knowledge
The neurobiology of decision-making and clinical expertise elevates anesthesiology to a high level.
reliability.

The consolidation of this knowledge allows for the creation of educational frameworks based on...
based on evidence, capable of standardizing behaviors, reducing operational variability and
To increase perioperative safety in different surgical and institutional settings.

**Methodological Challenges**

Scientific research related to neuroanesthesia in neurolysis surgical procedures for
The treatment of sequelae of leprosy faces methodological complexities that
limit the advancement of level 1 evidence. Leprosy is a disease with a distribution
heterogeneous geographic distribution strongly correlated with social determinants of poverty, the
which hinders private funding and the structuring of robust clinical research. The scarcity
The allocation of resources to the investigation of neglected pathologies results in studies with
reduced samples.

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google


**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

Another critical methodological challenge lies in the objective and standardized measurement of the effectiveness of Nerve decompression in the long-term postoperative period. Assessment of regeneration. Axonal damage depends on electromyography exams and skin sensitivity tests with... Monofilaments of esthesiometry and neuropathic pain scales that exhibit high variability. Interobserver reliability. The absence of specific biochemical biomarkers to quantify the degree. The difficulty in nerve regeneration makes it difficult to accurately compare anesthetic techniques. The design of double-blind masked clinical trials in anesthesia research. Regional anesthesia and decompressive surgery encounter technical barriers in execution and ethical limitations that are difficult to overcome. It is impossible to blind the anesthesiologist or the... surgeon regarding the blocking technique or surgical procedure performed on the operating table. operations. Although masking the assessor of postoperative outcomes is feasible and While mandatory, controlling for the placebo effect in peripheral neurolysis surgery raises debates. complex bioethics in academia.

Within the scope of studies focused on occupational health and cognitive performance of Anesthesiologist, self-reporting and historical recall biases compromise accuracy. from data collected through questionnaires. Research on the prevalence of Syndrome Burnout, work-related stress, and chronic fatigue suffer from the natural resistance of healthcare professionals are reluctant to admit psychological vulnerabilities due to corporate stigma. This systematic underreporting masks the true magnitude of the existing psychosocial risks. To overcome the methodological challenges in this interprofessional area, it is necessary to create... of multicenter research consortia focused on reconstructive surgery for endemic diseases and in Use of rigorous prospective observational methodologies. Integration of databases. Computerized global systems and the application of structural equation models help to control the multiple socioeconomic and clinical confounding variables of the patients studied. Investing in transparent and independent scientific methodologies is the way forward.

## Future Perspectives

The future of neuroanesthesia and surgical rehabilitation in leprosy will be marked by development and clinical application of ultrasound imaging systems High resolution and wearable miniaturized transducers. These advanced technologies This will allow mapping the microarchitecture of intraneural nerve fascicles with high resolution. Micrometric, identifying exact points of fibrotic constriction and precise anatomical guides.

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google
RCMOS – Multidisciplinary Scientific Journal The Knowledge.
ISSN: 2675-9128. São Paulo-SP.

**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

for the regional locking needle. This level of imaging will revolutionize the safety of procedures, eliminating blind spots.

Advances in tissue engineering and biomaterials will introduce the use of conductors. bioabsorbable materials for nerve regeneration and hydrogels enriched with factors Cell growth and stem cells at the site of peripheral neurolysis. The anesthesiologist will act. in the pharmacological modulation of the perineural microenvironment, administering adjuvants that prolong local inflammatory blockade and stimulate early intraneural angiogenesis in post-operative. This combined approach will accelerate the functional recovery time of the affected limbs.

In the field of occupational health, the introduction of continuous physiological monitoring systems, through smart wearable sensors integrated with artificial intelligence, it will allow To monitor the anesthesiologist's health in real time. These devices will monitor the heart rate, heart rate variability, biomechanical posture and Cognitive fatigue levels can be measured through vocal pattern analysis or eye tracking. during long surgical sessions. The system will issue preventive alerts or suggest breaks. organized automatic ergonomics.

Assisted robotics and surgical telerobotics will expand the reach of humanitarian actions and Specialized volunteering across continental distances. Experts located in large... Urban centers of excellence may guide or perform microsurgical procedures of Neurolysis in remote and isolated areas using high-precision robotic platforms. connected to ultra-high-speed internet networks. Local neuroanesthetic support will be optimized through tele-guidance systems in regional anesthesia, democratizing the access.

In the field of applied neuroscience, the development of brain-computer interfaces Non-invasive and repeated transcranial magnetic stimulation systems will open new avenues. to accelerate cortical plasticity and post-neurolysis motor rehabilitation. Mental training The patient's treatment, combined with central neuromodulatory stimulation, will help the brain recover. somatotopic maps lost due to chronic sensory deprivation resulting from leprosy.

The consolidation of these technologies will transform the management of chronic diseases.

## Conclusion

The complex clinical approach involving applied neuroanesthesia and procedures of

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google
RCMOS - Multidisciplinary Scientific Journal The Knowledge.
ISSN: 2675-9128. São Paulo-SP.

**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

Surgical decompressive rehabilitation in leprosy reinforces the need for a practice
A highly specialized doctor deeply committed to social responsibility.
Addressing the consequences of a neglected endemic disease requires professionals to overcome challenges.
of conventional service models and the fusion of advanced knowledge in
Neurology, pharmacology, and reconstructive microsurgery.
As reviewed scientific evidence demonstrates, the success of peripheral neurolysis in
Prevention and reversal of clawing and physical deformities intrinsically depend on precision.
intraoperative technique and maintenance of favorable microcirculatory conditions
Intraneural. Specialized neuroanesthetic management, based on regional anesthesia.
Guided by high-resolution ultrasound, it acts as a pillar of biosafety that
It protects neural tissues against secondary iatrogenic injuries.
The social and humanitarian impact resulting from the engagement of medical specialists in actions
Volunteering and highly productive assistance workshops reveal the transformative power
of medicine based on scientific solidarity. By restoring the ability to work and relieving
The suffering caused by chronic neuropathic pain, these humanitarian surgical interventions
They restore citizenship, dignity, and autonomy to historically underserved populations and
vulnerable.

From an occupational medicine perspective, the operating room environment presents challenges.
severe occupational hazards that must be continuously mapped and anticipated by management.
Hospital settings. Continuous exposure to residual anesthetic gases, ionizing radiation, and charges.
unfavorable ergonomic conditions, coupled with the severe psychosocial pressure of crisis management.
Perioperative care requires the implementation of robust preventive programs that ensure...
physical and mental health of the anesthesiologist.
The contributions of applied neuroscience to human clinical performance shed light on
the intricate neural processes of decision-making under acute stress in the operating room.
The adoption of resource management tools in medical crises and training.
Focused on maintaining situational awareness, they minimize the impact of stress on the...
cortical executive functions, ensuring cognitive stability and reducing the incidence of
Medical errors in critical settings.

The methodological challenges arising from the scarcity of resources allocated to the study of diseases.
Neglected tropical climates require the mobilization of the international academic community for the
Development of multicenter collaborative registries based on real-world data.
Overcoming sampling limitations and subjectivity in the assessment of neurological outcomes.

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Machine Translated by Google


**Year VI, v.1 2025 | Submission: 09/11/2025 | Accepted: 09/14/2025 | Publication: 09/17/2025**

Post-operative care is an urgent step towards consolidating robust global clinical guidelines and transparent in this area of operation.

More than contributing to the technical improvement of anesthesia in high-risk procedures Despite the complexity, the evidence analyzed demonstrates that the knowledge derived from Cognitive neuroscience, occupational health, and crisis management have the potential to... to be systematized into replicable professional training methodologies. Protocols focused on situational awareness, structured communication, emotional regulation, and Fatigue management and decision-making under pressure can be incorporated into the training. and to the continuing education of perioperative teams, strengthening the safety culture.

From this perspective, neuroanesthesia transcends its traditional role as an aid. clinical expertise and contributes to the development of high-level organizational models. reliability, capable of transforming scientific evidence into sustainable practices of Human performance, error prevention, and patient protection.

**References**

Brazilian Federal Council of Medicine. Occupational health standards and ergonomic risks in the surgical center environment for anesthesiologists. *Brazilian Journal of Occupational Medicine,* v. 41, n. 4, p. 202-215, 2024.

Brazilian Society of Neurosciences. Neurocognitive architecture and medical decision-making under acute stress in highly complex environments. *Journal of Applied Neuroscience and Health,* v. 18, n. 1, p. 55-68, 2023.

MINISTRY OF HEALTH (Brazil). *National guidelines for prevention, disability and surgical rehabilitation in leprosy.* Brasília: Ministry of Health, 2022.

OLIVEIRA, JLB et al. The impact of voluntary peripheral neurolysis workshops on the functional recovery of patients with chronic endemic diseases. *Annals of the Brazilian Academy of Humanitarian Medicine,* v. 12, n. 2, p. 114–126, 2025.

WORLD HEALTH ORGANIZATION (WHO). *Global leprosy (Hansen's disease) strategy 2021-2030: towards zero leprosy.* New Delhi: WHO Regional Office for South-East Asia, 2021.

This article is published in open access under the Creative Commons Attribution license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.