RCMOS – Revista Científica Multidisciplinar O Saber.
ISSN: 2675-9128. São Paulo - SP.
Year VII, v.1 2026 | Submission: 06/19/2026 | Accepted: 06/22/2026 | Publication: 06/25/2026

Female Weight Training: Comparative Analysis of Training Methodologies on Hypertrophy Parameters in Experienced Practitioners

Musculação Feminina: Análise Comparativa de Metodologias de Treino Sobre os Parâmetros de Hipertrofia em Praticantes Experientes

Musculación Femenina: Análisis Comparativo de Metodologías de Entrenamiento sobre los Parámetros de Hipertrofia en Practicantes Experimentadas

Emily Suelen Meireles Cabral – UNIFATECIE, emilymeeireles@gmail.com – Acadêmica de educação física
Mylena Mikaellen Beraldo – UNIFATECIE, mylena.beraldo@fatecie.edu.br - Orientadora. Professora do curso de educação física

Abstract:
This study aims to analyze and compare the effects of different resistance training methodologies on muscle hypertrophy parameters in experienced women. The research addresses traditional training and high metabolic volume training, discussing their characteristics, applications, and physiological impacts. The study is justified by the increasing participation of women in resistance training programs and the need for specific prescription strategies. The proposed methodology is characterized as quantitative, experimental, and field-based. The literature review was conducted using scientific articles, books, and academic documents. Expected results indicate that both methodologies can promote hypertrophy, albeit through distinct mechanisms. The study aims to contribute to scientific advancement and the professional practice of physical education professionals.

Keywords:
Weight training; Hypertrophy; Strength training; Women; Physical Education.

Resumo:
O presente estudo tem como objetivo analisar e comparar os efeitos de diferentes metodologias de treinamento de força sobre os parâmetros de hipertrofia muscular em mulheres experientes. A pesquisa aborda o treinamento tradicional e o treinamento de alto volume metabólico, discutindo suas características, aplicações e impactos fisiológicos. O estudo justifica-se pela crescente participação feminina em programas de musculação e pela necessidade de estratégias específicas de prescrição. A metodologia proposta caracteriza-se como quantitativa, experimental e de campo. A revisão bibliográfica foi desenvolvida a partir de artigos científicos, livros e documentos acadêmicos. Os resultados esperados indicam que ambas as metodologias podem promover hipertrofia, porém por mecanismos distintos. Espera-se contribuir para o avanço científico e para a atuação profissional do educador físico.

Palavras-chave:
Musculação; Hipertrofia; Treinamento de força; Mulheres; Educação Física.

Resumen:
Este estudio tiene como objetivo analizar y comparar los efectos de diferentes metodologías de entrenamiento de fuerza sobre los parámetros de hipertrofia muscular en mujeres experimentadas. La investigación aborda el entrenamiento tradicional y el entrenamiento de alto volumen metabólico, analizando sus características, aplicaciones e impactos fisiológicos. El estudio se justifica por la creciente participación femenina en los programas de musculación y por la necesidad de estrategias específicas de prescripción. La metodología propuesta se caracteriza por ser cuantitativa, experimental y de campo. La revisión bibliográfica se desarrolló

a partir de artículos científicos, libros y documentos académicos. Los resultados esperados indican que ambas metodologías pueden promover la hipertrofia, aunque mediante mecanismos diferentes. Se espera contribuir al avance científico y a la actuación profesional del educador físico.

Palabras clave:
Musculación; Hipertrofia; Entrenamiento de fuerza; Mujeres; Educación Física.

INTRODUCTION

Weight training has become one of the most common physical activities in gyms. The increased participation of women highlights the need for specific studies on the effects of strength training in women. Historically, most research has been conducted with men, leaving gaps in our understanding of women's adaptations to resistance training. Currently, it is known that women show positive responses to progressive overload, although there are physiological particularities that must be considered.

Muscle hypertrophy is one of the main goals for women who practice weight training. In this context, different training methodologies have been used to maximize results.

This study seeks to determine whether there are significant differences between traditional methodologies and high-volume metabolic methodologies in promoting muscle hypertrophy in experienced women.

The overall objective is to analyze and compare the effects of these methodologies on muscle hypertrophy parameters. Specific objectives include evaluating body composition, muscle thickness, maximum strength, and subjective perception of effort. The relevance of this work lies in its contribution to evidence-based professional practice.

DEVELOPMENT

Strength training consists of performing exercises against external resistance with the goal of producing neuromuscular adaptations capable of improving physical performance, health, and body composition. These adaptations involve structural, metabolic, and functional changes that result in increased muscle strength, localized muscle endurance, power, and hypertrophy.

The scientific principles of training include biological individuality, specificity, overload, continuity, and reversibility. These principles guide the development of training programs and ensure that the stimuli applied are capable of promoting positive adaptations.

RCMOS – Revista Científica Multidisciplinar O Saber.
ISSN: 2675-9128. São Paulo - SP.
Year VII, v.1 2026 | Submission: 06/19/2026 | Accepted: 06/22/2026 | Publication: 06/25/2026

Progressive overload is considered one of the most important factors for muscle hypertrophy, as it continuously stimulates muscles to adapt to increasing levels of physical demand. Without proper load progression, gains plateau and physiological adaptations decrease.

Mechanisms of Muscle Hypertrophy

Muscle hypertrophy is the result of complex cellular processes triggered by resistance training. Schoenfeld (2013) proposed three main mechanisms involved in this process: mechanical tension, metabolic stress, and muscle damage.

Mechanical tension is produced when a muscle generates force against significant resistance. This mechanism activates molecular pathways related to muscle growth, especially the mTOR pathway , considered fundamental for protein synthesis.

Metabolic stress occurs due to the accumulation of metabolites produced during intense exercise, such as lactate, hydrogen ions, and inorganic phosphate. This metabolic environment favors hormonal and cellular responses associated with muscle growth.

Muscle damage, in turn, refers to the micro-tears caused by exercise, especially during eccentric actions. Although it is not the primary hypertrophic mechanism, it plays an important role in muscle remodeling processes.

Specifics of Female Hypertrophy

Women have specific physiological characteristics that can influence their response to strength training. These characteristics include lower testosterone levels, a higher percentage of body fat, and differences in muscle fiber distribution.

Despite hormonal differences, studies show that women experience similar relative gains in strength and hypertrophy compared to men when subjected to appropriate training protocols. This is because the cellular mechanisms responsible for hypertrophy are largely shared between the sexes.

Another relevant aspect relates to the menstrual cycle, which can influence variables such as performance, recovery, and subjective perception of effort. However, the literature still presents conflicting results regarding the magnitude of these effects.

RCMOS – Revista Científica Multidisciplinar O Saber.
ISSN: 2675-9128. São Paulo - SP.
Year VII, v.1 2026 | Submission: 06/19/2026 | Accepted: 06/22/2026 | Publication: 06/25/2026

Traditional Training

Traditional training is characterized by the use of loads between 65% and 85% of one repetition maximum (1RM), generally performed in three to five sets with moderate recovery intervals.

This methodology prioritizes the systematic progression of load and allows for a high degree of control over training variables. Its main advantage lies in its ability to simultaneously develop muscle strength and hypertrophy.

Several studies demonstrate that traditional training promotes significant increases in muscle mass, especially when combined with gradual progression of loads and adequate protein intake.

High Metabolic Volume Training

High-volume metabolic training methods aim to maximize metabolic stress by increasing total workload and reducing recovery intervals.

Among the most commonly used techniques are:

- Drop-set;
- Rest -pause;
- Bi-set ;
- Tri-set ; ● Giant series;
- Partial repetitions.

These strategies prolong the time under stress and increase the recruitment of motor units, favoring hypertrophic adaptations.

Recent research indicates that high-volume protocols can produce results comparable to traditional methods in terms of hypertrophy, although they often generate greater fatigue and discomfort during training.

The drop set consists of performing a set to muscle failure followed by an immediate reduction in weight and continuation of the exercise without a rest interval. This technique significantly increases metabolic stress and muscle recruitment.

Its main advantage lies in its ability to prolong hypertrophic stimulation within a short period of time . However, overuse can increase fatigue and compromise muscle recovery.

RCMOS – Revista Científica Multidisciplinar O Saber.
ISSN: 2675-9128. São Paulo - SP.
Year VII, v.1 2026 | Submission: 06/19/2026 | Accepted: 06/22/2026 | Publication: 06/25/2026

rest -pause method is characterized by taking brief pauses during the exercise series. After reaching momentary fatigue, the practitioner takes a short rest and continues the exercise with the same load.

This strategy allows you to increase the total number of repetitions performed, raising the training volume without significantly reducing the intensity of the stimulus.

Hormonal Responses to Strength Training in Women

Hormonal responses play a fundamental role in the adaptations caused by resistance training. Although women have lower testosterone concentrations compared to men, this does not prevent the development of muscle hypertrophy. Studies indicate that hormones such as estrogen, growth hormone (GH), and insulin-like growth factor (IGF-1) actively participate in muscle recovery and growth processes.

Estrogen plays a protective role in muscle tissue, helping to reduce muscle damage and promoting recovery after intense training sessions. In addition, this hormone helps maintain bone mass and joint integrity, important aspects for women who perform high-intensity exercise.

Growth hormone levels temporarily increase after resistance training sessions, especially when using protocols with high workloads and short recovery intervals. This increase is associated with the stimulation of protein synthesis and the improvement of the body's regenerative processes.

Therefore, despite hormonal differences between the sexes, women have a physiological environment favorable to muscle development when subjected to properly structured programs.

Training Volume and Muscle Hypertrophy

Training volume refers to the total amount of work performed during a specific training session or period. It is generally calculated by multiplying the number of sets, repetitions, and weight used.

Several studies indicate that volume is one of the most important variables for muscle hypertrophy. Schoenfeld et al. (2019) observed that individuals subjected to higher training volumes showed greater gains in muscle mass when compared to groups that performed less work.

RCMOS – Revista Científica Multidisciplinar O Saber.
ISSN: 2675-9128. São Paulo - SP.
Year VII, v.1 2026 | Submission: 06/19/2026 | Accepted: 06/22/2026 | Publication: 06/25/2026

In experienced female athletes, gradually increasing training volume can be an important strategy to continue promoting muscle adaptations. However, excessive volumes can compromise recovery and increase the risk of cumulative fatigue.

Therefore, the professional must seek a balance between stimulation and recovery, adjusting the volume according to the individual capacity of the practitioner.

Training Intensity and Muscle Recruitment

Training intensity refers to the magnitude of the load used during exercise. Traditionally, it is believed that high loads are essential for the development of muscle hypertrophy.

However, recent studies demonstrate that moderate and even light loads can promote significant muscle growth when performed close to concentric failure. This occurs because progressive fatigue leads to the recruitment of high-threshold muscle fibers, responsible for force production and increased muscle mass.

Despite this, high loads remain important for developing maximum strength and maintaining neuromuscular efficiency. Therefore, programs that combine different intensity zones tend to produce more complete results.

Time Under Stress

Time under tension represents the period during which the muscle remains active during a series of exercises. This variable has received considerable attention in the scientific literature due to its potential influence on hypertrophic mechanisms.

Protocols that increase time under tension promote greater metabolic stress and prolong muscle activation. Methods such as drop sets, rest -pause, and controlled repetitions utilize this principle to enhance hypertrophic stimulation.

However, excessively increasing the time under tension can reduce the load used, compromising mechanical stress. Therefore, this variable must be manipulated in a balanced way within the training plan.

Muscle Recruitment and Adaptations to Training

RCMOS – Revista Científica Multidisciplinar O Saber.
ISSN: 2675-9128. São Paulo - SP.
Year VII, v.1 2026 | Submission: 06/19/2026 | Accepted: 06/22/2026 | Publication: 06/25/2026

Recovery is a fundamental component for the success of any training program. It is during rest periods that the physiological processes responsible for tissue repair and muscle growth occur.

Sleep deprivation, inadequate nutrition, and overtraining can significantly impair results. Studies show that individuals who sleep less than seven hours a night experience reduced recovery capacity and lower efficiency in muscle protein synthesis.

Furthermore, inadequate recovery can lead to overreaching and overtraining, conditions characterized by decreased performance and increased physical and mental fatigue.

Therefore, training planning should include appropriate recovery strategies to maximize hypertrophic results.

Nutrition and Muscle Hypertrophy in Women

Nutrition plays an indispensable role in the process of muscle hypertrophy. Strength training provides the necessary stimulus for muscle growth, but proper nutrition provides the substrates needed for protein synthesis.

Consuming high-quality protein is considered essential for muscle development. It is recommended that bodybuilders consume adequate amounts of protein throughout the day to optimize anabolic processes.

In addition to proteins, carbohydrates and fats play important roles related to energy supply and hormone production. Nutritional deficiencies can significantly limit the results obtained from training.

Therefore, the integration between training and nutrition is a determining factor for the success of programs aimed at female muscle hypertrophy.

Recent Scientific Evidence Regarding Female Hypertrophy

Research published in recent years has shown that trained women exhibit a high capacity for adaptation to resistance training. The results indicate that both traditional protocols and high-volume methodologies can promote significant increases in muscle mass.

However, the literature indicates that individual response plays an important role in the observed results. Genetic, hormonal, nutritional, and behavioral factors directly influence adaptations to training.

RCMOS – Revista Científica Multidisciplinar O Saber.
ISSN: 2675-9128. São Paulo - SP.
Year VII, v.1 2026 | Submission: 06/19/2026 | Accepted: 06/22/2026 | Publication: 06/25/2026

Given this, the importance of individualizing exercise programs grows, allowing each participant to receive stimuli compatible with their specific needs and goals.

Current evidence reinforces the idea that there is no single ideal methodology for all women. The success of training depends on the proper combination of volume, intensity, frequency, recovery, and adherence to the program.

Comparison Between Methodologies

The comparison between traditional training and high-volume metabolic training is one of the most discussed topics in exercise science. While traditional training emphasizes mechanical stress, high-volume methods prioritize metabolic stress.

Current evidence suggests that both mechanisms are capable of promoting muscle hypertrophy. However, there is still no absolute consensus on which strategy produces greater gains in trained women.

Some studies indicate that higher volumes may promote muscle growth in experienced individuals. Other authors argue that the intensity of the load remains the determining variable for superior adaptations.

Therefore, a balanced combination of intensity and volume seems to represent the most appropriate strategy for optimizing hypertrophic results.

Practical Applications for Physical Education Professionals

Research findings on female hypertrophy have direct implications for professional practice. The choice of methodology should consider factors such as training experience, individual goals, time availability, recovery capacity, and adherence to the program.

Physical education professionals should avoid prescriptions based solely on market trends or empirical experiences, prioritizing approaches grounded in scientific evidence.

Furthermore, it is essential to monitor indicators such as weekly volume, intensity, subjective perception of effort, and muscle recovery to ensure positive adaptations and minimize the risk of overtraining.

Scientific literature demonstrates that resistance training promotes significant adaptations in the musculoskeletal system. Muscle hypertrophy occurs in response to a combination of mechanical tension, metabolic stress, and cellular processes related to protein

synthesis. Several authors point out that the proper manipulation of training variables, including volume, intensity, frequency, and recovery intervals, directly influences the results obtained.

In the context of women, there is a growing interest in training programs aimed at increasing muscle mass. Recent studies indicate that women can achieve similar relative gains to men when subjected to appropriate protocols. However, hormonal factors, anthropometric characteristics, and differences related to recovery should be considered during prescription.

In addition to aesthetic benefits, weight training contributes to bone health, improved functional capacity, prevention of chronic diseases, and increased quality of life. Therefore, regular strength training is an important strategy for promoting health.

Comparing different training methodologies allows us to understand which strategies offer the best balance between effectiveness, safety, and adherence. Physical education professionals should base their decisions on up-to-date scientific evidence, respecting the principles of biological individuality and load progression.

FINAL CONSIDERATIONS

In conclusion, female bodybuilding represents a field of growing scientific and professional interest. Both traditional training and high-volume metabolic methodologies have the potential to promote muscle hypertrophy. The choice of the most appropriate strategy depends on individual goals, training experience, and the practitioner's recovery capacity.

Analysis of the scientific literature demonstrates that female muscle hypertrophy is a multifactorial phenomenon influenced by variables related to training, nutrition, recovery, and individual characteristics.

Both traditional training and high-volume metabolic methods have the potential to promote significant muscle mass gains in experienced women. However, the adaptations observed depend on the proper manipulation of training variables and the individualization of programs.

The reviewed studies suggest that there is no universally superior methodology for all practitioners. Instead, the strategic combination of different methods appears to represent the most efficient alternative for maximizing hypertrophic results.

In conclusion, strength training is an extremely effective tool for developing muscle hypertrophy in women, and it is essential that its prescription be based on up-to-date scientific evidence and adapted to the individual needs of each participant.

The expected results of this research may contribute to the development of more efficient and safer training programs, expanding scientific knowledge related to women's adaptations to resistance training. Further studies are recommended to deepen the understanding of women's physiological responses to different training methodologies.

REFERENCES

BOMPA, TO; BUZZICHELLI, C. **Periodization of sports training.** São Paulo: Phorte , 2019.

FLECK, SJ; KRAEMER, WJ **Fundamentals of strength training** . Porto Alegre: Artmed, 2017.

GENTIL, P. et al. **Comparison of muscle hypertrophy responses in males and females .** The Journal of Strength & Conditioning Research , vol. 31, no. 2, p. 517-522, 2017.

HUNTER, SK **Sex differences in humans Fatigability mechanisms and insight to physiological responses.** Acta Physiologica , v. 210, no. 4, p. 768-789, 2016.

ROBERGS, RA et al. **Gender differences in muscle strength and muscle mass .** Sports Medicine, vol. 48, n. 1, p. 47-57, 2018.

SCHOENFELD, BJ **Potential mechanisms for a role of metabolic stress in hypertrophic adaptations to resistance training.** Sports Medicine, vol. 43, no. 3, p. 179-194, 2013.

SCHOENFELD, BJ et al. **Resistance training volume enhancers muscle hypertrophy but not strength in trained men .** Medicine & Science in Sports & Exercise , vol. 51, n. 1, p. 94-103, 2019